

# DRINFEL'D DOUBLES AND LUSZTIG'S SYMMETRIES OF TWO-PARAMETER QUANTUM GROUPS

NANTEL BERGERON<sup>a, \*</sup>, YUN GAO<sup>a, \*\*</sup>, NAIHONG HU<sup>b, †</sup>

<sup>a</sup>Department of Mathematics and Statistics, York University  
Toronto, Ontario, Canada M3J 1P3

<sup>b</sup>Department of Mathematics, East China Normal University  
Shanghai 200062, People's Republic of China

First Version on Sept. 1, 2004, Revised Version on April 15, 2005

**ABSTRACT.** We find the defining structures of two-parameter quantum groups  $U_{r,s}(\mathfrak{g})$  corresponding to the orthogonal and the symplectic Lie algebras, which are realized as Drinfel'd doubles. We further investigate the environment conditions upon which the Lusztig's symmetries exist between  $(U_{r,s}(\mathfrak{g}), \langle, \rangle)$  and its associated object  $(U_{s^{-1}, r^{-1}}(\mathfrak{g}), \langle | \rangle)$ .

## 0. INTRODUCTION

In the early 1990's, many authors investigated two-parameter or multiparameter quantum groups. These authors include Kulish [14], Reshetikhin [18], Sudbery [21], Takeuchi [22], Artin, Schelter and Tate [1], Du, Parshall and Wang [9], Dobrev and Parashar [7], Jing [11], Chin and Musson [6], etc. (for details, see the introduction of [3] and references therein). Their works focused on quantized function algebras and quantum enveloping algebras only for type  $A$  cases. In 2001, Benkart and Witherspoon [3], motivated by the work on generalizations of algebras generated by the down and up operators on posets (see down-up algebras in [2]), obtained the structure of two-parameter quantum enveloping algebras corresponding to the general linear Lie algebra  $\mathfrak{gl}_n$  and the special linear Lie algebra  $\mathfrak{sl}_n$  (which was also studied earlier by Takeuchi in [22] with a different motivation). These two-parameter quantum enveloping algebras of type  $A$  were proved to

---

\*Supported in part by the NSERC and the CRC of Canada

\*\*Supported in part by the NSERC of Canada

†Supported by the NNSF (Grant 10431040), the TRAPOYT and the SFUDP from the MOE of China, the Shanghai Priority Academic Discipline from the SEC.

1991 *Mathematics Subject Classification.* 17B37, 81R50.

*Key words and phrases.* 2-parameter quantum group, Hopf (skew-)dual pairing, Drinfel'd double, Lusztig's symmetry.

have Drinfel'd doubles structures. Furthermore, in [4] they studied the finite-dimensional weight representation theory under the assumption that  $rs^{-1}$  is not a root of unity, and showed that the finite-dimensional weight modules are completely reducible. This is analogous to the classical semisimple Lie algebras  $\mathfrak{g}$  and their one-parameter quantum groups  $U_q(\mathfrak{g})$  of Drinfel'd-Jimbo type (with  $q$  generic). Since then, a systematic study for the two-parameter quantum group of type  $A$  has been further developed by Benkart and Witherspoon, etc. (see [5] and references therein). However, up to now, how to find the defining relations of two-parameter quantum groups of other types was still open and an interesting task.

In the present paper, we give the definitions of two-parameter quantum groups of types  $B$ ,  $C$ ,  $D$ , which are new to our knowledge. We also prove that these quantum groups we have constructed have a Drinfel'd doubles structures. We then investigate the Lusztig's symmetry properties for these two-parameter quantum groups for  $B$ ,  $C$  and  $D$  types as well as for type  $A$ . Particularly, we derive some interesting  $(r, s)$ -identities held in  $U_{r,s}(\mathfrak{g})$  (see Lemma 3.6).

The paper is organized as follows. In Section 1, we present the definitions of the two-parameter quantum groups corresponding to the orthogonal Lie algebras  $\mathfrak{so}_{2n+1}$  or  $\mathfrak{so}_{2n}$  and the symplectic Lie algebras  $\mathfrak{sp}_{2n}$ , together with the Hopf algebra structure. As it was done for type  $A$  case in [3], we prove in Section 2 that two-parameter quantum groups  $U_{r,s}(\mathfrak{g})$  are characterized as Drinfel'd doubles  $\mathcal{D}(\mathcal{B}, \mathcal{B}')$  of the Hopf subalgebras  $\mathcal{B}$  (upper part) and  $\mathcal{B}'$  (lower part) with respect to a skew-dual paring. As a by-product of the double structure, one naturally gets a standard triangular decomposition of  $U_{r,s}(\mathfrak{g})$  (see Corollary 2.6), which plays a crucial role in finite dimensional weight representation theory of  $U_{r,s}(\mathfrak{g})$ . The double structure also leads to a direct approach to the triangular decomposition of quantum groups  $U_q(\mathfrak{g})$  of Drinfel'd-Jimbo type, which, originally, was obtained nontrivially by Lusztig [16] and Rosso [20] in different ways. Section 3 is devoted to investigating the Lusztig's symmetries for the two-parameter quantum groups. A striking feature of these symmetries is that they exist as  $\mathbb{Q}$ -isomorphisms between  $U_{r,s}(\mathfrak{g})$  and the associated object  $U_{s^{-1}, r^{-1}}(\mathfrak{g})$  only when  $\text{rank}(\mathfrak{g}) = 2$ . When  $\text{rank}(\mathfrak{g}) > 2$ , the sufficient and necessary condition for the existence of Lusztig's symmetries between  $U_{r,s}(\mathfrak{g})$  and its associated object forces  $U_{r,s}(\mathfrak{g})$  to take the "one-parameter" form  $U_{q,q^{-1}}(\mathfrak{g})$  where  $r = s^{-1} = q$ . In other words, we prove that when  $\text{rank}(\mathfrak{g}) > 2$ , the Lusztig's symmetries exist only for the one-parameter quantum groups  $U_{q,q^{-1}}(\mathfrak{g})$  as  $\mathbb{Q}(q)$ -automorphisms (rather than merely  $\mathbb{Q}$ -isomorphisms). In this case, these symmetries coincide, modulo identification, with the usual Lusztig symmetries on quantum groups  $U_q(\mathfrak{g})$  of Drinfel'd-Jimbo type (see [12, 13, 15]). Some necessary calculation data are collected in Section 4, which are useful to the proof of Proposition 2.3.

## 1. TWO-PARAMETER QUANTUM GROUPS OF TYPES $B$ , $C$ , $D$

Let  $\mathbb{K} = \mathbb{Q}(r, s)$  denote a field of rational functions with two indeterminates  $r$ ,  $s$ , or a subfield of  $\mathbb{C}$  with two-parameters  $r$ ,  $s$  with assumption  $r^2 \neq s^2$ .

In order to reveal more about the defining structure of the “two-parameter” quantum groups, we present the definitions case by case. In particular, the  $(r, s)$ -Serre relations for the  $D_n$  case have some special features for the vertices  $n-1$  and  $n$ , which degenerate into the usual commutative relations for the one-parameter case (i.e., when  $rs = 1$  in (D5) below).

(I) Assume  $\Psi$  is a finite root system of type  $B_n$  with  $\Pi$  a base of simple roots. Regard  $\Psi$  as a subset of a Euclidean space  $E = \mathbb{R}^n$  with an inner product  $(\cdot, \cdot)$ . Let  $\epsilon_1, \dots, \epsilon_n$  denote an orthonormal basis of  $E$ , and suppose  $\Pi = \{\alpha_i = \epsilon_i - \epsilon_{i+1} \mid 1 \leq i < n\} \cup \{\alpha_n = \epsilon_n\}$  and  $\Psi = \{\pm\epsilon_i \pm \epsilon_j \mid 1 \leq i \neq j \leq n\} \cup \{\pm\epsilon_i \mid 1 \leq i \leq n\}$ . In this case, set  $r_i = r^{(\alpha_i, \alpha_i)}$  and  $s_i = s^{(\alpha_i, \alpha_i)}$  so that  $r_1 = \dots = r_{n-1} = r^2$ ,  $r_n = r$  and  $s_1 = \dots = s_{n-1} = s^2$ ,  $s_n = s$ .

Let  $U = U_{r,s}(\mathfrak{so}_{2n+1})$  be the unital associative algebra over  $\mathbb{Q}(r, s)$  generated by  $e_i, f_i, \omega_i^{\pm 1}, \omega'_i{}^{\pm 1}$  ( $1 \leq i \leq n$ ), subject to the following relations (B1)–(B7):

(B1) The  $\omega_i^{\pm 1}, \omega'_j{}^{\pm 1}$  all commute with one another and  $\omega_i \omega_i^{-1} = \omega'_j \omega'_j{}^{-1} = 1$ .

(B2) For  $1 \leq i \leq n$  and  $1 \leq j < n$ , we have

$$\begin{aligned} \omega_j e_i \omega_j^{-1} &= r_j^{(\epsilon_j, \alpha_i)} s_j^{(\epsilon_{j+1}, \alpha_i)} e_i, & \omega_j f_i \omega_j^{-1} &= r_j^{-(\epsilon_j, \alpha_i)} s_j^{-(\epsilon_{j+1}, \alpha_i)} f_i, \\ \omega_n e_j \omega_n^{-1} &= r_n^{2(\epsilon_n, \alpha_j)} e_j, & \omega_n f_j \omega_n^{-1} &= r_n^{-2(\epsilon_n, \alpha_j)} f_j, \\ \omega_n e_n \omega_n^{-1} &= r_n^{(\epsilon_n, \alpha_n)} s_n^{-(\epsilon_n, \alpha_n)} e_n, & \omega_n f_n \omega_n^{-1} &= r_n^{-(\epsilon_n, \alpha_n)} s_n^{(\epsilon_n, \alpha_n)} f_n. \end{aligned}$$

(B3) For  $1 \leq i \leq n$  and  $1 \leq j < n$ , we have

$$\begin{aligned} \omega'_j e_i \omega'_j{}^{-1} &= s_j^{(\epsilon_j, \alpha_i)} r_j^{(\epsilon_{j+1}, \alpha_i)} e_i, & \omega'_j f_i \omega'_j{}^{-1} &= s_j^{-(\epsilon_j, \alpha_i)} r_j^{-(\epsilon_{j+1}, \alpha_i)} f_i, \\ \omega'_n e_j \omega'_n{}^{-1} &= s_n^{2(\epsilon_n, \alpha_j)} e_j, & \omega'_n f_j \omega'_n{}^{-1} &= s_n^{-2(\epsilon_n, \alpha_j)} f_j, \\ \omega'_n e_n \omega'_n{}^{-1} &= s_n^{(\epsilon_n, \alpha_n)} r_n^{-(\epsilon_n, \alpha_n)} e_n, & \omega'_n f_n \omega'_n{}^{-1} &= s_n^{-(\epsilon_n, \alpha_n)} r_n^{(\epsilon_n, \alpha_n)} f_n. \end{aligned}$$

(B4) For  $1 \leq i, j \leq n$ , we have

$$[e_i, f_j] = \delta_{ij} \frac{\omega_i - \omega'_i}{r_i - s_i}.$$

(B5) For any  $i, j$  with  $|i - j| > 1$ , we have the  $(r, s)$ -Serre relations:

$$[e_i, e_j] = [f_i, f_j] = 0.$$

(B6) For  $1 \leq i < n$ ,  $1 \leq j < n-1$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} e_i^2 e_{i+1} - (r_i + s_i) e_i e_{i+1} e_i + (r_i s_i) e_{i+1} e_i^2 &= 0, \\ e_{j+1}^2 e_j - (r_{j+1}^{-1} + s_{j+1}^{-1}) e_{j+1} e_j e_{j+1} + (r_{j+1}^{-1} s_{j+1}^{-1}) e_j e_{j+1}^2 &= 0, \\ e_n^3 e_{n-1} - (r_n^{-2} + r_n^{-1} s_n^{-1} + s_n^{-2}) e_n^2 e_{n-1} e_n + (r_n^{-1} s_n^{-1}) (r_n^{-2} + r_n^{-1} s_n^{-1} + s_n^{-2}) e_n e_{n-1} e_n^2 \\ - (r_n^{-3} s_n^{-3}) e_{n-1} e_n^3 &= 0. \end{aligned}$$

(B7) For  $1 \leq i < n$ ,  $1 \leq j < n-1$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} f_{i+1}f_i^2 - (r_i+s_i)f_i f_{i+1}f_i + (r_i s_i)f_i^2 f_{i+1} &= 0, \\ f_j f_{j+1}^2 - (r_{j+1}^{-1}+s_{j+1}^{-1})f_{j+1}f_j f_{j+1} + (r_{j+1}^{-1}s_{j+1}^{-1})f_{j+1}^2 f_j &= 0, \\ f_{n-1}f_n^3 - (r_n^{-2}+r_n^{-1}s_n^{-1}+s_n^{-2})f_n f_{n-1}f_n^2 + (r_n^{-1}s_n^{-1})(r_n^{-2}+r_n^{-1}s_n^{-1}+s_n^{-2})f_n^2 f_{n-1}f_n \\ &\quad - (r_n^{-3}s_n^{-3})f_n^3 f_{n-1} = 0. \end{aligned}$$

(II) Assume  $\Psi$  is a finite root system of type  $C_n$  with  $\Pi$  a base of simple roots. Regard  $\Psi$  as a subset of a Euclidean space  $E = \mathbb{R}^n$  with an inner product  $(\cdot, \cdot)$ . Let  $\epsilon_1, \dots, \epsilon_n$  denote an orthonormal basis of  $E$ , and suppose  $\Pi = \{\alpha_i = \epsilon_i - \epsilon_{i+1} \mid 1 \leq i < n\} \cup \{\alpha_n = 2\epsilon_n\}$  and  $\Psi = \{\pm\epsilon_i \pm \epsilon_j \mid 1 \leq i \neq j \leq n\} \cup \{2\epsilon_i \mid 1 \leq i \leq n\}$ . In this case, set  $r_i = r^{\frac{(\alpha_j, \alpha_i)}{2}}$  and  $s_i = s^{\frac{(\alpha_i, \alpha_i)}{2}}$  so that  $r_1 = \dots = r_{n-1} = r$ ,  $r_n = r^2$  and  $s_1 = \dots = s_{n-1} = s$ ,  $s_n = s^2$ .

Let  $U = U_{r,s}(\mathfrak{sp}_{2n})$  be the unital associative algebra over  $\mathbb{Q}(r, s)$  generated by  $e_i, f_i, \omega_i^{\pm 1}, \omega'_i{}^{\pm 1}$  ( $1 \leq i \leq n$ ), subject to the following relations (C1)–(C7):

(C1) The  $\omega_i^{\pm 1}, \omega'_j{}^{\pm 1}$  all commute with one another and  $\omega_i \omega_i^{-1} = \omega'_j \omega'_j{}^{-1} = 1$ .

(C2) For  $1 \leq i \leq n$  and  $1 \leq j < n$ , we have

$$\begin{aligned} \omega_j e_i \omega_j^{-1} &= r^{(\epsilon_j, \alpha_i)} s^{(\epsilon_{j+1}, \alpha_i)} e_i, & \omega_j f_i \omega_j^{-1} &= r^{-(\epsilon_j, \alpha_i)} s^{-(\epsilon_{j+1}, \alpha_i)} f_i, \\ \omega_n e_j \omega_n^{-1} &= r^{2(\epsilon_n, \alpha_j)} e_j, & \omega_n f_j \omega_n^{-1} &= r^{-2(\epsilon_n, \alpha_j)} f_j, \\ \omega_n e_n \omega_n^{-1} &= r^{(\epsilon_n, \alpha_n)} s^{-(\epsilon_n, \alpha_n)} e_n, & \omega_n f_n \omega_n^{-1} &= r^{-(\epsilon_n, \alpha_n)} s^{(\epsilon_n, \alpha_n)} f_n. \end{aligned}$$

(C3) For  $1 \leq i \leq n$  and  $1 \leq j < n$ , we have

$$\begin{aligned} \omega'_j e_i \omega'_j{}^{-1} &= s^{(\epsilon_j, \alpha_i)} r^{(\epsilon_{j+1}, \alpha_i)} e_i, & \omega'_j f_i \omega'_j{}^{-1} &= s^{-(\epsilon_j, \alpha_i)} r^{-(\epsilon_{j+1}, \alpha_i)} f_i, \\ \omega'_n e_j \omega'_n{}^{-1} &= s^{2(\epsilon_n, \alpha_j)} e_j, & \omega'_n f_j \omega'_n{}^{-1} &= s^{-2(\epsilon_n, \alpha_j)} f_j, \\ \omega'_n e_n \omega'_n{}^{-1} &= s^{(\epsilon_n, \alpha_n)} r^{-(\epsilon_n, \alpha_n)} e_n, & \omega'_n f_n \omega'_n{}^{-1} &= s^{-(\epsilon_n, \alpha_n)} r^{(\epsilon_n, \alpha_n)} f_n. \end{aligned}$$

(C4) For  $1 \leq i, j \leq n$ , we have

$$[e_i, f_j] = \delta_{ij} \frac{\omega_i - \omega'_i}{r_i - s_i}.$$

(C5) For any  $i, j$  with  $|i - j| > 1$ , we have the  $(r, s)$ -Serre relations:

$$[e_i, e_j] = [f_i, f_j] = 0.$$

(C6) For  $1 \leq i < n-1$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} e_i^2 e_{i+1} - (r+s) e_i e_{i+1} e_i + (rs) e_{i+1} e_i^2 &= 0, \\ e_{n-1}^3 e_n - (r^2+rs+s^2) e_{n-1}^2 e_n e_{n-1} + (rs) (r^2+rs+s^2) e_{n-1} e_n e_{n-1}^2 - (rs)^3 e_n e_{n-1}^3 &= 0, \\ e_{i+1}^2 e_i - (r^{-1}+s^{-1}) e_{i+1} e_i e_{i+1} + (r^{-1}s^{-1}) e_i e_{i+1}^2 &= 0, \\ e_n^2 e_{n-1} - (r_n^{-1}+s_n^{-1}) e_n e_{n-1} e_n + (r_n^{-1}s_n^{-1}) e_{n-1} e_n^2 &= 0. \end{aligned}$$

(C7) For  $1 \leq i < n-1$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} f_{i+1}f_i^2 - (r+s)f_if_{i+1}f_i + (rs)f_i^2f_{i+1} &= 0, \\ f_nf_{n-1}^3 - (r^2+rs+s^2)f_{n-1}f_nf_{n-1}^2 + (rs)(r^2+rs+s^2)f_{n-1}^2f_nf_{n-1} - (rs)^3f_{n-1}^3f_n &= 0, \\ f_if_{i+1}^2 - (r^{-1}+s^{-1})f_{i+1}f_if_{i+1} + (r^{-1}s^{-1})f_{i+1}^2f_i &= 0, \\ f_{n-1}f_n^2 - (r_n^{-1}+s_n^{-1})f_nf_{n-1}f_n + (r_n^{-1}s_n^{-1})f_n^2f_{n-1} &= 0. \end{aligned}$$

(III) Assume  $\Psi$  is a finite root system of type  $D_n$  with  $\Pi$  a base of simple roots. Regard  $\Psi$  as a subset of a Euclidean space  $E = \mathbb{R}^n$  with an inner product  $(\cdot, \cdot)$ . Let  $\epsilon_1, \dots, \epsilon_n$  denote an orthonormal basis of  $E$ , and suppose  $\Pi = \{\alpha_i = \epsilon_i - \epsilon_{i+1} \mid 1 \leq i < n\} \cup \{\alpha_n = \epsilon_{n-1} + \epsilon_n\}$  and  $\Psi = \{\pm\epsilon_i \pm \epsilon_j \mid 1 \leq i \neq j \leq n\}$ . In this case, set  $r_i = r^{\frac{(\alpha_i, \alpha_i)}{2}}$  and  $s_i = s^{\frac{(\alpha_i, \alpha_i)}{2}}$  so that  $r_1 = \dots = r_n = r$  and  $s_1 = \dots = s_n = s$ .

Let  $U = U_{r,s}(\mathfrak{so}_{2n})$  be the unital associative algebra over  $\mathbb{Q}(r, s)$  generated by  $e_i, f_i, \omega_i^{\pm 1}, \omega'_i{}^{\pm 1}$  ( $1 \leq i \leq n$ ), subject to the following relations (D1)–(D7):

(D1) The  $\omega_i^{\pm 1}, \omega'_j{}^{\pm 1}$  all commute with one another and  $\omega_i\omega_i^{-1} = \omega'_j\omega'_j{}^{-1} = 1$ .

(D2) For  $1 \leq i \leq n$ ,  $1 \leq j < n$ , and  $1 \leq k (\neq n-1) \leq n$ , we have

$$\begin{aligned} \omega_j e_i \omega_j^{-1} &= r^{(\epsilon_j, \alpha_i)} s^{(\epsilon_{j+1}, \alpha_i)} e_i, & \omega_j f_i \omega_j^{-1} &= r^{-(\epsilon_j, \alpha_i)} s^{-(\epsilon_{j+1}, \alpha_i)} f_i, \\ \omega_n e_k \omega_n^{-1} &= r^{(\epsilon_{n-1}, \alpha_k)} s^{-(\epsilon_n, \alpha_k)} e_k, & \omega_n f_k \omega_n^{-1} &= r^{-(\epsilon_{n-1}, \alpha_k)} s^{(\epsilon_n, \alpha_k)} f_k, \\ \omega_n e_{n-1} \omega_n^{-1} &= r^{(\epsilon_n, \alpha_{n-1})} s^{-(\epsilon_{n-1}, \alpha_{n-1})} e_{n-1}, & \omega_n f_{n-1} \omega_n^{-1} &= r^{-(\epsilon_n, \alpha_{n-1})} s^{(\epsilon_{n-1}, \alpha_{n-1})} f_{n-1}. \end{aligned}$$

(D3) For  $1 \leq i \leq n$ ,  $1 \leq j < n$ , and  $1 \leq k (\neq n-1) \leq n$ , we have

$$\begin{aligned} \omega'_j e_i \omega'_j{}^{-1} &= s^{(\epsilon_j, \alpha_i)} r^{(\epsilon_{j+1}, \alpha_i)} e_i, & \omega'_j f_i \omega'_j{}^{-1} &= s^{-(\epsilon_j, \alpha_i)} r^{-(\epsilon_{j+1}, \alpha_i)} f_i, \\ \omega'_n e_k \omega'_n{}^{-1} &= s^{(\epsilon_{n-1}, \alpha_k)} r^{-(\epsilon_n, \alpha_k)} e_k, & \omega'_n f_k \omega'_n{}^{-1} &= s^{-(\epsilon_{n-1}, \alpha_k)} r^{(\epsilon_n, \alpha_k)} f_k, \\ \omega'_n e_{n-1} \omega'_n{}^{-1} &= s^{(\epsilon_n, \alpha_{n-1})} r^{-(\epsilon_{n-1}, \alpha_{n-1})} e_{n-1}, & \omega'_n f_{n-1} \omega'_n{}^{-1} &= s^{-(\epsilon_n, \alpha_{n-1})} r^{(\epsilon_{n-1}, \alpha_{n-1})} f_{n-1}. \end{aligned}$$

(D4) For  $1 \leq i, j \leq n$ , we have

$$[e_i, f_j] = \delta_{ij} \frac{\omega_i - \omega'_i}{r - s}.$$

(D5) For any  $1 \leq i \neq j \leq n$  but  $(i, j) \notin \{(n-1, n), (n, n-1)\}$  with  $a_{ij} = 0$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} [e_i, e_j] &= [f_i, f_j] = 0, \\ e_{n-1}e_n &= rs e_n e_{n-1}, & f_n f_{n-1} &= rs f_{n-1} f_n. \end{aligned}$$

(D6) For  $1 \leq i < j \leq n$  with  $a_{ij} = -1$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} e_i^2 e_j - (r+s) e_i e_j e_i + (rs) e_j e_i^2 &= 0, \\ e_j^2 e_i - (r^{-1}+s^{-1}) e_j e_i e_j + (r^{-1}s^{-1}) e_i e_j^2 &= 0. \end{aligned}$$

(D7) For  $1 \leq i < j \leq n$  with  $a_{ij} = -1$ , we have the  $(r, s)$ -Serre relations:

$$\begin{aligned} f_j f_i^2 - (r+s) f_i f_j f_i + (rs) f_i^2 f_j &= 0, \\ f_i f_j^2 - (r^{-1} + s^{-1}) f_j f_i f_j + (r^{-1} s^{-1}) f_j^2 f_i &= 0. \end{aligned}$$

In summary, let  $U = U_{r,s}(\mathfrak{g})$  for  $\mathfrak{g} = \mathfrak{so}_{2n+1}$ ,  $\mathfrak{so}_{2n}$  and  $\mathfrak{sp}_{2n}$  denote the two-parameter quantum orthogonal groups and quantum symplectic groups, respectively.

The following fact is straightforward to check.

**Proposition 1.1.** *The algebra  $U_{r,s}(\mathfrak{g})$  ( $\mathfrak{g} = \mathfrak{so}_{2n+1}$ ,  $\mathfrak{so}_{2n}$ , or  $\mathfrak{sp}_{2n}$ ) is a Hopf algebra under the comultiplication, the counit and the antipode below:*

$$\begin{aligned} \Delta(\omega_i^{\pm 1}) &= \omega_i^{\pm 1} \otimes \omega_i^{\pm 1}, & \Delta(\omega_i'^{\pm 1}) &= \omega_i'^{\pm 1} \otimes \omega_i'^{\pm 1}, \\ \Delta(e_i) &= e_i \otimes 1 + \omega_i \otimes e_i, & \Delta(f_i) &= 1 \otimes f_i + f_i \otimes \omega_i', \\ \varepsilon(\omega_i^{\pm 1}) &= \varepsilon(\omega_i'^{\pm 1}) = 1, & \varepsilon(e_i) &= \varepsilon(f_i) = 0, \\ S(\omega_i^{\pm 1}) &= \omega_i^{\mp 1}, & S(\omega_i'^{\pm 1}) &= \omega_i'^{\mp 1}, \\ S(e_i) &= -\omega_i^{-1} e_i, & S(f_i) &= -f_i \omega_i'^{-1}. \end{aligned}$$

**Remarks.** (1) When  $r = q$  and  $s = q^{-1}$ , the Hopf algebra  $U_{q,q^{-1}}(\mathfrak{g})$  modulo the Hopf ideal generated by the elements  $\omega_i' - \omega_i^{-1}$  ( $1 \leq i \leq n$ ), is the one-parameter quantum groups  $U_q(\mathfrak{g})$  of Drinfel'd-Jimbo type.

(2) As usual, we define respectively the left-adjoint and the right-adjoint action in the Hopf algebra  $U_{r,s}(\mathfrak{g})$  as

$$\mathrm{ad}_l a(b) = \sum_{(a)} a_{(1)} b S(a_{(2)}), \quad \mathrm{ad}_r a(b) = \sum_{(a)} S(a_{(1)}) b a_{(2)},$$

where  $\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}$ , for any  $a, b \in U_{r,s}(\mathfrak{g})$ .

Using adjoint actions, the  $(r, s)$ -Serre relations (X5), (X6) and (X7) (here  $X = B, C, D$ ) in  $U_{r,s}(\mathfrak{g})$  become simply:

$$(\mathrm{ad}_l e_i)^{1-a_{ij}}(e_j) = 0, \quad \text{for any } i \neq j, \quad (\text{X6})$$

$$(\mathrm{ad}_r f_i)^{1-a_{ij}}(f_j) = 0, \quad \text{for any } i \neq j. \quad (\text{X7})$$

According to the data on the prime root systems of the classical simple Lie algebras, the following basic lemma is clear. This will play a crucial role in ensuring the compatibility of the defining relations (X2) & (X3) for  $U_{r,s}(\mathfrak{g})$ , especially, in the proof of Theorem 2.5.

**Lemma 1.2.** *For the prime root systems of the Lie algebras  $\mathfrak{g} = \mathfrak{sl}_n$ ,  $\mathfrak{so}_{2n+1}$ ,  $\mathfrak{so}_{2n}$ , and  $\mathfrak{sp}_{2n}$ , the following identities hold:*

$$\begin{aligned} (\epsilon_{j+1}, \alpha_i) &= -(\epsilon_i, \alpha_j), & (i, j < n), & \text{for any } \mathfrak{g}, \\ (\epsilon_{j+1}, \alpha_n) &= \begin{cases} -(\epsilon_n, \alpha_j), & (j < n), \\ -2(\epsilon_n, \alpha_j), & (j < n), \end{cases} & \text{for } \mathfrak{g} = \mathfrak{so}_{2n+1}, \\ & & \text{for } \mathfrak{g} = \mathfrak{sp}_{2n}, \\ (\epsilon_j, \alpha_n) &= \begin{cases} -(\epsilon_n, \alpha_{j-1}), & (j \leq n, j \neq n-1), \\ (\epsilon_{n-1}, \alpha_{n-1}), & (j = n-1) \end{cases} & \text{for } \mathfrak{g} = \mathfrak{so}_{2n}. \end{aligned}$$

## 2. DRINFELD'S QUANTUM DOUBLES AND ROSSO FORM

**Definition 2.1.** A (Hopf) dual pairing of two Hopf algebras  $\mathcal{A}$  and  $\mathcal{U}$  (see [12, 3.2.1], or [13, 1.2.5]) is a bilinear form  $\langle \cdot, \cdot \rangle : \mathcal{U} \times \mathcal{A} \longrightarrow \mathbb{K}$  such that

$$\langle f, 1_{\mathcal{A}} \rangle = \varepsilon_{\mathcal{U}}(f), \quad \langle 1_{\mathcal{U}}, a \rangle = \varepsilon_{\mathcal{A}}(a), \quad (1)$$

$$\langle f, a_1 a_2 \rangle = \langle \Delta_{\mathcal{U}}(f), a_1 \otimes a_2 \rangle, \quad \langle f_1 f_2, a \rangle = \langle f_1 \otimes f_2, \Delta_{\mathcal{A}}(a) \rangle, \quad (2)$$

for all  $f, f_1, f_2 \in \mathcal{U}$ , and  $a, a_1, a_2 \in \mathcal{A}$ . Here  $\varepsilon_{\mathcal{U}}$  and  $\varepsilon_{\mathcal{A}}$  denote the counits of  $\mathcal{U}$  and  $\mathcal{A}$ , respectively, and  $\Delta_{\mathcal{U}}$  and  $\Delta_{\mathcal{A}}$  are their comultiplications.

A direct consequence of the defining properties above is that the dual pairing satisfies

$$\langle S_{\mathcal{U}}(f), a \rangle = \langle f, S_{\mathcal{A}}(a) \rangle, \quad f \in \mathcal{U}, \quad a \in \mathcal{A},$$

where  $S_{\mathcal{U}}, S_{\mathcal{A}}$  denote the respective antipodes of  $\mathcal{U}$  and  $\mathcal{A}$ .

**Definition 2.2.** A bilinear form  $\langle \cdot, \cdot \rangle : \mathcal{U} \times \mathcal{A} \longrightarrow \mathbb{K}$  is called a skew-dual pairing of two Hopf algebras  $\mathcal{A}$  and  $\mathcal{U}$  (see [13, 8.2.1]) if  $\langle \cdot, \cdot \rangle : \mathcal{U}^{\text{cop}} \times \mathcal{A} \longrightarrow \mathbb{K}$  is a dual pairing of the two Hopf algebras  $\mathcal{A}$  and  $\mathcal{U}^{\text{cop}}$ . Here  $\mathcal{U}^{\text{cop}}$  is the Hopf algebra having the opposite comultiplication to the Hopf algebra  $\mathcal{U}$  and  $S_{\mathcal{U}^{\text{cop}}} = S_{\mathcal{U}}^{-1}$  if  $S_{\mathcal{U}}$  is invertible.

Let  $\mathcal{B} = B(\mathfrak{g})$  (resp.  $\mathcal{B}' = B'(\mathfrak{g})$ ) denote the Hopf subalgebra of  $U = U_{r,s}(\mathfrak{g})$  generated by  $e_j, \omega_j^{\pm 1}$  (resp.  $f_j, \omega_j^{\pm 1}$ ) with  $1 \leq j < n$  for  $\mathfrak{g} = \mathfrak{sl}_n$ , and with  $1 \leq j \leq n$  for  $\mathfrak{g} = \mathfrak{so}_{2n+1}$ ,  $\mathfrak{so}_{2n}$ , or  $\mathfrak{sp}_{2n}$ . Note that the case  $\mathfrak{g} = \mathfrak{sl}_n$  was done in [3]. The main ideas of this section comes from this work. Since our statement is based on Definition 2.2 and slightly different, we also include the case  $\mathfrak{g} = \mathfrak{sl}_n$  in Proposition 2.3 and Theorem 2.5 below.

**Proposition 2.3.** There exists a unique skew-dual pairing  $\langle \cdot, \cdot \rangle : B'(\mathfrak{g}) \times B(\mathfrak{g}) \longrightarrow \mathbb{Q}(r, s)$  of the Hopf subalgebras  $B(\mathfrak{g})$  and  $B'(\mathfrak{g})$ , for  $\mathfrak{g} = \mathfrak{sl}_n, \mathfrak{so}_{2n+1}, \mathfrak{so}_{2n}$ , or  $\mathfrak{sp}_{2n}$  such that

$$\langle f_i, e_j \rangle = \delta_{ij} \frac{1}{s_i - r_i}, \quad \text{for any } \mathfrak{g}, \quad (3)$$

$$\langle \omega'_i, \omega_j \rangle = r^{(\epsilon_j, \alpha_i)} s^{(\epsilon_{j+1}, \alpha_i)}, \quad i < n, \quad j < n, \quad \text{for } \mathfrak{sl}_n, \quad (4A)$$

$$\langle \omega'_i, \omega_j \rangle = \begin{cases} r^{2(\epsilon_j, \alpha_i)} s^{2(\epsilon_{j+1}, \alpha_i)}, & i \leq n, \quad j < n, \\ r^{2(\epsilon_n, \alpha_i)}, & i < n, \quad j = n, \\ r^{(\epsilon_n, \alpha_n)} s^{-(\epsilon_n, \alpha_n)}, & i = j = n. \end{cases} \quad \text{for } \mathfrak{so}_{2n+1}, \quad (4B)$$

$$\langle \omega'_i, \omega_j \rangle = \begin{cases} r^{(\epsilon_j, \alpha_i)} s^{(\epsilon_{j+1}, \alpha_i)}, & i \leq n, \quad j < n, \\ r^{2(\epsilon_n, \alpha_i)}, & i < n, \quad j = n, \\ r^{(\epsilon_n, \alpha_n)} s^{-(\epsilon_n, \alpha_n)}, & i = j = n. \end{cases} \quad \text{for } \mathfrak{sp}_{2n}, \quad (4C)$$

$$\langle \omega'_i, \omega_j \rangle = \begin{cases} r^{(\epsilon_j, \alpha_i)} s^{(\epsilon_{j+1}, \alpha_i)}, & i \leq n, \quad j < n, \\ r^{(\epsilon_{n-1}, \alpha_i)} s^{-(\epsilon_n, \alpha_i)}, & i \neq n-1, \quad j = n, \\ r^{(\epsilon_n, \alpha_{n-1})} s^{-(\epsilon_{n-1}, \alpha_{n-1})}, & i = n-1, \quad j = n. \end{cases} \quad \text{for } \mathfrak{so}_{2n}, \quad (4D)$$

$$\langle \omega_i^{\pm 1}, \omega_j^{-1} \rangle = \langle \omega_i^{\pm 1}, \omega_j \rangle^{-1} = \langle \omega'_i, \omega_j \rangle^{\mp 1}, \quad \text{for any } \mathfrak{g}, \quad (5)$$

and all other pairs of generators are 0. Moreover, we have  $\langle S(a), S(b) \rangle = \langle a, b \rangle$  for  $a \in \mathcal{B}'$ ,  $b \in \mathcal{B}$ .

*Proof.* The uniqueness assertion is clear, since any skew-dual pairing of bialgebras is determined by the values on the generators. We proceed to prove the existence of the pairing.

We begin by defining on the generators a bilinear form  $\langle \cdot, \cdot \rangle : \mathcal{B}'^{\text{cop}} \times \mathcal{B} \longrightarrow \mathbb{Q}(r, s)$  satisfying (3), (4X) and (5). We then extend it to a bilinear form on  $\mathcal{B}'^{\text{cop}} \times \mathcal{B}$  by requiring that (1) and (2) hold for  $\Delta_{\mathcal{B}'^{\text{cop}}} = \Delta_{\mathcal{B}'}^{\text{op}}$ . We verify that the relations in  $\mathcal{B}$  and  $\mathcal{B}'$  are preserved, ensuring that the form is well-defined and is a dual pairing of  $\mathcal{B}$  and  $\mathcal{B}'^{\text{cop}}$ .

It is straightforward to check that the bilinear form preserves all the relations among the  $\omega_i^{\pm 1}$  in  $\mathcal{B}$  and the  $\omega'_i{}^{\pm 1}$  in  $\mathcal{B}'$ . We notice that Lemma 1.2 ensures the compatibility of the form defined above with the relations of (X2) and (X3) in  $\mathcal{B}$  and  $\mathcal{B}'$  respectively. This follows by definition (from (3), (4X), & (5)). We are left to verify that the form preserves the  $(r, s)$ -Serre relations in  $\mathcal{B}$  and  $\mathcal{B}'$ . It suffices to show that the form on  $\mathcal{B}'^{\text{cop}} \times \mathcal{B}$  preserves the  $(r, s)$ -Serre relations in  $\mathcal{B}$ , because the verification for  $\mathcal{B}'^{\text{cop}}$  is similar. To this end, we observe that the relations involving indices  $1 \leq i, j < n$  belong to  $\mathcal{B} \cap U_{r,s}(\mathfrak{sl}_n) = B(\mathfrak{sl}_n)$ . This has been checked by Benkart and Witherspoon in the type  $A$  case (see [3, Lemma 2.2]). We have thus reduced the proof to the rank 2 type  $B_2$  and  $C_2$ , and the rank 4 type  $D_4$ .

(I) Cases  $\mathfrak{g} = \mathfrak{sp}_{2n}$  and  $\mathfrak{so}_{2n+1}$ : we only need to consider the case  $n = 2$  ( $B_2$  and  $C_2$ ).

First, let us show that the form preserves the  $(r, s)$ -Serre relation of degree 3 in  $\mathcal{B}$ :

$$\langle X, e_1^3 e_2 - (r^2 + rs + s^2) e_1^2 e_2 e_1 + (rs)(r^2 + rs + s^2) e_1 e_2 e_1^2 - (rs)^3 e_2 e_1^3 \rangle = 0, \quad (C_2)$$

$$\begin{aligned} \langle X, e_2^3 e_1 - (r^{-2} + r^{-1} s^{-1} + s^{-2}) e_2^2 e_1 e_2 + (r^{-1} s^{-1})(r^{-2} + r^{-1} s^{-1} + s^{-2}) e_2 e_1 e_2^2 \\ - (r^{-1} s^{-1})^3 e_1 e_2^3 \rangle = 0, \end{aligned} \quad (B_2)$$

where  $X$  is any word in the generators of  $\mathcal{B}'$ . By definition, the left-hand side of each of the above identities respectively equals

$$\begin{aligned} \langle \Delta^{(3)}(X), e_1 \otimes e_1 \otimes e_1 \otimes e_2 - (r^2 + rs + s^2) e_1 \otimes e_1 \otimes e_2 \otimes e_1 \\ + (rs)(r^2 + rs + s^2) e_1 \otimes e_2 \otimes e_1 \otimes e_1 - (rs)^3 e_2 \otimes e_1 \otimes e_1 \otimes e_1 \rangle, \end{aligned} \quad (6C)$$

$$\begin{aligned} \langle \Delta^{(3)}(X), e_2 \otimes e_2 \otimes e_2 \otimes e_1 - (r^{-2} + r^{-1} s^{-1} + s^{-2}) e_2 \otimes e_2 \otimes e_1 \otimes e_2 \\ + (r^{-1} s^{-1})(r^{-2} + r^{-1} s^{-1} + s^{-2}) e_2 \otimes e_1 \otimes e_2 \otimes e_2 - (r^{-3} s^{-3}) e_1 \otimes e_2 \otimes e_2 \otimes e_2 \rangle, \end{aligned} \quad (6B)$$

where the  $\Delta$  corresponds to  $\Delta_{\mathcal{B}'}^{\text{op}}$ . In order for any one of these terms to be nonzero,  $X$  must involve exactly three  $f_1$  factors, one  $f_2$  factor, and arbitrarily many  $\omega'_j{}^{\pm 1}$  factors ( $j = 1, 2$ ).

We first consider the following four key cases:

(i) For type  $C_2$  and  $X = f_1^3 f_2$ , we have

$$\begin{aligned} \Delta^{(3)}(X) = (\omega'_1 \otimes \omega'_1 \otimes \omega'_1 \otimes f_1 + \omega'_1 \otimes \omega'_1 \otimes f_1 \otimes 1 + \omega'_1 \otimes f_1 \otimes 1 \otimes 1 + f_1 \otimes 1 \otimes 1 \otimes 1)^3 \\ \cdot (\omega'_2 \otimes \omega'_2 \otimes \omega'_2 \otimes f_2 + \omega'_2 \otimes \omega'_2 \otimes f_2 \otimes 1 + \omega'_2 \otimes f_2 \otimes 1 \otimes 1 + f_2 \otimes 1 \otimes 1 \otimes 1). \end{aligned}$$

In Appendix (4.1) we have listed only the terms in the expansion of  $\Delta^{(3)}(X)$  that have a non-zero contribution in (6C). Consequently, by properties (1) & (2) of  $\langle, \rangle$  on  $\mathcal{B}'^{\text{cop}} \times \mathcal{B}$ , the pairing (6C) becomes

$$\begin{aligned} & \langle f_1, e_1 \rangle^3 \langle f_2, e_2 \rangle (1 + 2\langle \omega'_1, \omega_1 \rangle + 2\langle \omega'_1, \omega_1 \rangle^2 + \langle \omega'_1, \omega_1 \rangle^3) \cdot \\ & \cdot (1 - r^2 \langle \omega'_1, \omega_2 \rangle) (1 - rs \langle \omega'_1, \omega_2 \rangle) (1 - s^2 \langle \omega'_1, \omega_2 \rangle) \\ & = 0, \quad (\text{since } \langle \omega'_1, \omega_2 \rangle = r^{-2}). \end{aligned}$$

For type  $B_2$ , when  $X = f_2^3 f_1$ , we only need to interchange the indices 1 and 2, and substitute  $r, s$  by  $r^{-1}, s^{-1}$  in the above identity, to obtain the pairing (6B) as

$$\begin{aligned} & \langle f_1, e_1 \rangle \langle f_2, e_2 \rangle^3 (1 + 2\langle \omega'_2, \omega_2 \rangle + 2\langle \omega'_2, \omega_2 \rangle^2 + \langle \omega'_2, \omega_2 \rangle^3) \\ & \cdot (1 - r^{-2} \langle \omega'_2, \omega_1 \rangle) (1 - r^{-1} s^{-1} \langle \omega'_2, \omega_1 \rangle) (1 - s^{-2} \langle \omega'_2, \omega_1 \rangle) \\ & = 0, \quad (\text{since } \langle \omega'_2, \omega_1 \rangle = s^2). \end{aligned}$$

(ii) For type  $C_2$  and  $X = f_2 f_1^3$ , the relevant terms of  $\Delta^{(3)}(X)$  for (6C) are listed in Appendix (4.2). The result of (6C) is then

$$\begin{aligned} & \langle f_1, e_1 \rangle^3 \langle f_2, e_2 \rangle (1 + 2\langle \omega'_1, \omega_1 \rangle + 2\langle \omega'_1, \omega_1 \rangle^2 + \langle \omega'_1, \omega_1 \rangle^3) \cdot \\ & \cdot (\langle \omega'_2, \omega_1 \rangle - r^2) (\langle \omega'_2, \omega_1 \rangle - rs) (\langle \omega'_2, \omega_1 \rangle - s^2) \\ & = 0, \quad (\text{since } \langle \omega'_2, \omega_1 \rangle = s^2). \end{aligned}$$

Again for type  $B_2$ , when  $X = f_1 f_2^3$  we need to interchange the indices 1 and 2, and substitute  $r, s$  by  $r^{-1}, s^{-1}$  in the above identity, to obtain the pairing (6B) as

$$\begin{aligned} & \langle f_1, e_1 \rangle \langle f_2, e_2 \rangle^3 (1 + 2\langle \omega'_2, \omega_2 \rangle + 2\langle \omega'_2, \omega_2 \rangle^2 + \langle \omega'_2, \omega_2 \rangle^3) \\ & \cdot (\langle \omega'_1, \omega_2 \rangle - r^{-2}) (\langle \omega'_1, \omega_2 \rangle - r^{-1} s^{-1}) (\langle \omega'_1, \omega_2 \rangle - s^{-2}) \\ & = 0, \quad (\text{since } \langle \omega'_1, \omega_2 \rangle = r^{-2}). \end{aligned}$$

(iii) For type  $C_2$  and  $X = f_1^2 f_2 f_1$ , the relevant terms of  $\Delta^{(3)}(X)$  for (6C) are listed in Appendix (4.3), and (6C) becomes

$$\begin{aligned} & \langle f_1, e_1 \rangle^3 \langle f_2, e_2 \rangle \left\{ (1 + 2\langle \omega'_1, \omega_1 \rangle + 2\langle \omega'_1, \omega_1 \rangle^2 + \langle \omega'_1, \omega_1 \rangle^3) (\langle \omega'_2, \omega_1 \rangle - (rs)^3 \langle \omega'_1, \omega_2 \rangle^2) \right. \\ & + (r^2 + rs + s^2) \left[ (rs) \langle \omega'_1, \omega_2 \rangle (1 + 2\langle \omega'_1, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle^2) \right. \\ & \quad + \langle \omega'_1, \omega_1 \rangle^2 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle^3 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle) \\ & \quad - (1 + \langle \omega'_1, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle \\ & \quad \left. \left. + 2\langle \omega'_1, \omega_1 \rangle^2 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle^3 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle) \right] \right\} \\ & = 0, \quad (\text{since } \langle \omega'_1, \omega_1 \rangle = rs^{-1}, \langle \omega'_1, \omega_2 \rangle = r^{-2}, \langle \omega'_2, \omega_1 \rangle = s^2). \end{aligned}$$

Similarly, for  $B_2$  and  $X = f_2^2 f_1 f_2$ , interchanging indices 1 and 2, and substituting  $r, s$  by  $r^{-1}, s^{-1}$  in the above identity, we obtain that the pairing (6B) is zero.

(iv) For type  $C_2$  and  $X = f_1 f_2 f_1^2$ , the relevant terms of  $\Delta^{(3)}(X)$  for (6C) are listed in Appendix (4.4), and (6C) becomes

$$\begin{aligned} & \langle f_1, e_1 \rangle^3 \langle f_2, e_2 \rangle (1 + 2\langle \omega'_1, \omega_1 \rangle + 2\langle \omega'_1, \omega_1 \rangle^2 + \langle \omega'_1, \omega_1 \rangle^3) (\langle \omega'_2, \omega_1 \rangle^2 - (rs)^3 \langle \omega'_1, \omega_2 \rangle) \\ & + (r^2 + rs + s^2) [ (rs) (1 + \langle \omega'_1, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle \\ & \quad + 2\langle \omega'_1, \omega_1 \rangle^2 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle^3 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle) \\ & \quad - \langle \omega'_2, \omega_1 \rangle (1 + 2\langle \omega'_1, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle^2 \\ & \quad + \langle \omega'_1, \omega_1 \rangle^2 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle + \langle \omega'_1, \omega_1 \rangle^3 \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle) ] \\ & = 0, \quad (\text{since } \langle \omega'_1, \omega_1 \rangle = rs^{-1}, \langle \omega'_1, \omega_2 \rangle = r^{-2}, \langle \omega'_2, \omega_1 \rangle = s^2). \end{aligned}$$

Again for type  $B_2$  and  $X = f_2 f_1 f_2^2$ , interchanging indices 1 and 2, and substituting  $r, s$  by  $r^{-1}, s^{-1}$  in the above identity, we obtain that the pairing (6B) is also zero.

Finally, we note the fact that if  $X = \omega'_2{}^p f_1^3 \omega'_1{}^q f_2 \omega'_j{}^s$  (resp.  $\omega'_1{}^q f_2^3 \omega'_2{}^p f_1 \omega'_j{}^s$ ), then (6C) (resp. (6B)) is  $\langle \omega'_2, \omega_1 \rangle^p \langle \omega'_1, \omega_2 \rangle^q$  times the corresponding quantity for  $X = f_1^3 f_2$  for  $C_2$  (resp.  $X = f_2^3 f_1$  for  $B_2$ ). That means, if  $X$  is any word involving exactly three  $f_1$  factors, one  $f_2$  factor, and arbitrarily many factors of  $\omega'_j{}^{\pm 1}$  ( $j = 1, 2$ ), then (6C) (resp. (6B)) is just a scalar multiple of one of the quantities we have already calculated, and therefore equal 0.

Next, we verify that the  $(r, s)$ -Serre relation of degree 2 in  $\mathcal{B}$  are preserved by the form. We do so by showing the identities

$$\langle X, e_2^2 e_1 - (r^{-2} + s^{-2}) e_2 e_1 e_2 + (r^{-2} s^{-2}) e_1 e_2^2 \rangle = 0, \quad (7C)$$

$$\langle X, e_1^2 e_2 - (r^2 + s^2) e_1 e_2 e_1 + (r^2 s^2) e_2 e_1^2 \rangle = 0, \quad (7B)$$

where  $X$  is any word in the generators of  $\mathcal{B}'$ . It is enough to consider three monomials:  $X = f_2^2 f_1, f_2 f_1 f_2$  and  $f_1 f_2^2$  for  $C_2$  (resp.  $X = f_1^2 f_2, f_1 f_2 f_1$  and  $f_2 f_1^2$  for  $B_2$ ).

(i) For type  $C_2$  and  $X = f_2^2 f_1$ , the relevant terms of  $\Delta^{(2)}(X)$  are listed in Appendix (4.5), and the left-hand side of (7C) equals

$$\langle f_2, e_2 \rangle^2 \langle f_1, e_1 \rangle (1 + \langle \omega'_2, \omega_2 \rangle) (1 - r^{-2} \langle \omega'_2, \omega_1 \rangle) (1 - s^{-2} \langle \omega'_2, \omega_1 \rangle),$$

which is 0, since  $\langle \omega'_2, \omega_1 \rangle = s^2$ .

For  $B_2$  and  $X = f_1^2 f_2$ , the left-hand side of (7B) is

$$\langle f_1, e_1 \rangle^2 \langle f_2, e_2 \rangle (1 + \langle \omega'_1, \omega_1 \rangle) (1 - r^2 \langle \omega'_1, \omega_2 \rangle) (1 - s^2 \langle \omega'_1, \omega_2 \rangle),$$

which is 0, since  $\langle \omega'_1, \omega_2 \rangle = r^{-2}$ .

(ii) Similarly, for type  $C_2$  and  $X = f_1 f_2^2$  (resp. type  $B_2$  and  $X = f_2 f_1^2$ ), the left-hand side of (7C) and (7B) respectively are equal to

$$\begin{aligned} & \langle f_1, e_1 \rangle \langle f_2, e_2 \rangle^2 (1 + \langle \omega'_2, \omega_2 \rangle) (\langle \omega'_1, \omega_2 \rangle - r^{-2}) (\langle \omega'_1, \omega_2 \rangle - s^{-2}) = 0, \quad (\text{as } \langle \omega'_1, \omega_2 \rangle = r^{-2}); \\ & \langle f_1, e_1 \rangle^2 \langle f_2, e_2 \rangle (1 + \langle \omega'_1, \omega_1 \rangle) (\langle \omega'_2, \omega_1 \rangle - r^2) (\langle \omega'_2, \omega_1 \rangle - s^2) = 0, \quad (\text{as } \langle \omega'_2, \omega_1 \rangle = s^2). \end{aligned}$$

(iii) For type  $C_2$  and  $X = f_2 f_1 f_2$ , the relevant terms of  $\Delta^{(2)}(X)$  are listed in Appendix (4.6), and the left-hand side of (7C) equals

$$\begin{aligned} & \langle f_2, e_2 \rangle^2 \langle f_1, e_1 \rangle \{ (1 + \langle \omega'_2, \omega_2 \rangle) (\langle \omega'_1, \omega_2 \rangle + r^{-2} s^{-2} \langle \omega'_2, \omega_1 \rangle) \\ & - (r^{-2} + s^{-2}) (1 + \langle \omega'_2, \omega_2 \rangle \langle \omega'_1, \omega_2 \rangle \langle \omega'_2, \omega_1 \rangle) \}, \end{aligned}$$

which is equal to 0, since  $\langle \omega'_2, \omega_2 \rangle = r^2 s^{-2}$ ,  $\langle \omega'_1, \omega_2 \rangle = r^{-2}$  and  $\langle \omega'_2, \omega_1 \rangle = s^2$ .

Similarly, for  $B_2$  and  $X = f_1 f_2 f_1$ , the left-hand side of (7B) is

$$\begin{aligned} & \langle f_1, e_1 \rangle^2 \langle f_2, e_2 \rangle \{ (1 + \langle \omega'_1, \omega_1 \rangle) (\langle \omega'_2, \omega_1 \rangle + r^2 s^2 \langle \omega'_1, \omega_2 \rangle) \\ & - (r^2 + s^2) (1 + \langle \omega'_1, \omega_1 \rangle \langle \omega'_2, \omega_1 \rangle \langle \omega'_1, \omega_2 \rangle) \}, \end{aligned}$$

which is equal to 0, since  $\langle \omega'_1, \omega_1 \rangle = r^2 s^{-2}$ ,  $\langle \omega'_1, \omega_2 \rangle = r^{-2}$  and  $\langle \omega'_2, \omega_1 \rangle = s^2$ .

A similar process shows that the relations in  $\mathcal{B}'^{\text{cop}}$  are preserved in the case when  $\mathfrak{g} = \mathfrak{sp}_{2n}$ , or  $\mathfrak{so}_{2n+1}$ .

(II) Case  $\mathfrak{g} = \mathfrak{so}_{2n}$ : we only need to consider the type  $D_4$ .

For  $j = 3$  or  $4$ , we let  $\mathcal{B}_j$  (resp.  $\mathcal{B}'_j$ ) denote the subalgebra generated by  $\omega_i, e_i$  (resp.  $\omega'_i, f_i$ ) with  $i \in \{1, 2, j\}$ . Note that  $\mathcal{B}_3 \cong \mathcal{B}_4 \cong B(\mathfrak{sl}_4)$  and  $\mathcal{B}_3 \cap \mathcal{B}_4 = B(\mathfrak{sl}_3) \subset B(\mathfrak{sl}_4)$ ;  $\mathcal{B}'_3 \cong \mathcal{B}'_4 \cong B'(\mathfrak{sl}_4)$  and  $\mathcal{B}'_3 \cap \mathcal{B}'_4 = B'(\mathfrak{sl}_3) \subset B'(\mathfrak{sl}_4)$ . The existence of pairings on  $\mathcal{B}'^{\text{cop}}_j \times \mathcal{B}_j$  ( $j = 3, 4$ ), whose restrictions on  $B'(\mathfrak{sl}_3)^{\text{cop}} \times B(\mathfrak{sl}_3)$  are consistent, has been proved in [3, Lemma 2.2]. Therefore, in order to show that the form defined on  $\mathcal{B}'^{\text{cop}} \times \mathcal{B}$  preserves the  $(r, s)$ -Serre relations in  $\mathcal{B}$ , it remains to verify that the  $(r, s)$ -Serre relation in  $\mathcal{B}$  involving indices 3, 4 in (D5) is preserved. We need only to consider two cases:  $X = f_i f_j$  for  $(i, j) = (3, 4)$ , or  $(4, 3)$ .

$$\begin{aligned} & \langle X, e_3 e_4 - (rs) e_4 e_3 \rangle \\ &= \langle f_i \omega'_j \otimes f_j + \omega'_i f_j \otimes f_i, e_3 \otimes e_4 - (rs) e_4 \otimes e_3 \rangle \\ &= \langle f_i, e_3 \rangle \langle f_j, e_4 \rangle - (rs) \langle f_i, e_4 \rangle \langle f_j, e_3 \rangle \\ &\quad + \langle \omega'_i, \omega_3 \rangle \langle f_j, e_3 \rangle \langle f_i, e_4 \rangle - (rs) \langle \omega'_i, \omega_4 \rangle \langle f_j, e_4 \rangle \langle f_i, e_3 \rangle \\ &= \begin{cases} \frac{1}{(s-r)^2} (1 - (rs) \langle \omega'_3, \omega_4 \rangle), & (i, j) = (3, 4), \\ \frac{1}{(s-r)^2} (\langle \omega'_4, \omega_3 \rangle - (rs)), & (i, j) = (4, 3). \end{cases} \\ &= 0, \quad (\text{since } \langle \omega'_3, \omega_4 \rangle = r^{-1} s^{-1}, \text{ \& } \langle \omega'_4, \omega_3 \rangle = rs). \end{aligned}$$

Similarly, we can prove that the  $(r, s)$ -Serre relation (D5) is preserved for  $\mathcal{B}'^{\text{cop}}$ .

(I) and (II) together complete the proof of the Proposition.  $\square$

**Definition 2.4.** For any two Hopf algebras  $\mathcal{A}$  and  $\mathcal{U}$  paired by a skew-dual pairing  $\langle \cdot, \cdot \rangle$ , one may form the Drinfel'd (quantum) double  $\mathcal{D}(\mathcal{A}, \mathcal{U})$  as in [12, 3.2] or [13, 8.2]. This is a Hopf algebra whose underlying vector space is  $\mathcal{A} \otimes \mathcal{U}$  with the tensor product coalgebra structure and the algebra structure defined by

$$(a \otimes f)(a' \otimes f') = \sum \langle Su(f_{(1)}), a'_{(1)} \rangle \langle f_{(3)}, a'_{(3)} \rangle a a'_{(2)} \otimes f_{(2)} f', \quad (8)$$

for  $a, a' \in \mathcal{A}$  and  $f, f' \in \mathcal{U}$ , and whose antipode  $S$  is given by

$$S(a \otimes f) = (1 \otimes S_{\mathcal{U}}(f))(S_{\mathcal{A}}(a) \otimes 1). \quad (9)$$

Clearly, both mappings  $\mathcal{A} \ni a \mapsto a \otimes 1 \in \mathcal{D}(\mathcal{A}, \mathcal{U})$  and  $\mathcal{U} \ni f \mapsto 1 \otimes f \in \mathcal{D}(\mathcal{A}, \mathcal{U})$  are injective Hopf algebra homomorphisms. Let us denote the image  $a \otimes 1$  (resp.  $1 \otimes f$ ) of  $a$  (resp.  $f$ ) in  $\mathcal{D}(\mathcal{A}, \mathcal{U})$  by  $\hat{a}$  (resp.  $\hat{f}$ ). By (8), we have the following cross commutation relations between elements  $\hat{a}$  (for  $a \in \mathcal{A}$ ) and  $\hat{f}$  (for  $f \in \mathcal{U}$ ) in the algebra  $\mathcal{D}(\mathcal{A}, \mathcal{U})$ :

$$\hat{f} \hat{a} = \sum \langle S_{\mathcal{U}}(f_{(1)}), a_{(1)} \rangle \langle f_{(3)}, a_{(3)} \rangle \hat{a}_{(2)} \hat{f}_{(2)}, \quad (10)$$

$$\sum \langle f_{(1)}, a_{(1)} \rangle \hat{f}_{(2)} \hat{a}_{(2)} = \sum \hat{a}_{(1)} \hat{f}_{(1)} \langle f_{(2)}, a_{(2)} \rangle. \quad (11)$$

In fact, as an algebra, the double  $\mathcal{D}(\mathcal{A}, \mathcal{U})$  is the universal algebra generated by the algebras  $\mathcal{A}$  and  $\mathcal{U}$  with cross relation (10) or equivalently (11).

**Theorem 2.5.** *The two-parameter quantum group  $U = U_{r,s}(\mathfrak{g})$  is isomorphic to the Drinfel'd quantum double  $\mathcal{D}(\mathcal{B}, \mathcal{B}')$ , for  $\mathfrak{g} = \mathfrak{sl}_n, \mathfrak{so}_{2n+1}, \mathfrak{so}_{2n}$ , or  $\mathfrak{sp}_{2n}$ .*

*Proof.* Define a mapping  $\varphi : \mathcal{D}(\mathcal{B}, \mathcal{B}') \longrightarrow U_{r,s}(\mathfrak{g})$  by

$$\begin{aligned} \varphi(\hat{\omega}_i^{\pm 1}) &= \omega_i^{\pm 1}, & \varphi(\hat{\omega}_i'^{\pm 1}) &= \omega_i'^{\pm 1}, \\ \varphi(\hat{e}_i) &= e_i, & \varphi(\hat{f}_i) &= f_i. \end{aligned}$$

Note that by definition,  $\varphi$  preserves the coalgebra structures, the relations in  $\mathcal{B}$ , and the relations in  $\mathcal{B}'$ .

We next verify that the cross relations in the double  $\mathcal{D}(\mathcal{B}, \mathcal{B}')$  correspond to those in  $U$ .

By (11) and applying the comultiplication given in Proposition 1.1, we obtain the cross relations

$$\begin{aligned} \langle \omega_j'^{\pm 1}, \omega_i^{\pm 1} \rangle \hat{\omega}_j'^{\pm 1} \hat{\omega}_i^{\pm 1} &= \hat{\omega}_i^{\pm 1} \hat{\omega}_j'^{\pm 1} \langle \omega_j'^{\pm 1}, \omega_i^{\pm 1} \rangle, \\ \langle \omega_j', \omega_i \rangle \hat{\omega}_j' \hat{e}_i &= \hat{e}_i \hat{\omega}_j' \langle \omega_j', 1 \rangle, \\ \langle 1, \omega_j^{-1} \rangle \hat{f}_i \hat{\omega}_j^{-1} &= \hat{\omega}_j^{-1} \hat{f}_i \langle \omega_i', \omega_j^{-1} \rangle, \\ \langle f_j, e_i \rangle \hat{\omega}_j' + \langle 1, \omega_i \rangle \hat{f}_j \hat{e}_i &= \hat{e}_i \hat{f}_j \langle \omega_j', 1 \rangle + \hat{\omega}_i \langle f_j, e_i \rangle. \end{aligned}$$

That is,

$$\begin{aligned} \hat{\omega}_j'^{\pm 1} \hat{\omega}_i^{\pm 1} &= \hat{\omega}_i^{\pm 1} \hat{\omega}_j'^{\pm 1}, \\ \hat{\omega}_j' \hat{e}_i \hat{\omega}_j'^{-1} &= \hat{e}_i \langle \omega_j', \omega_i \rangle^{-1}, \\ \hat{\omega}_j \hat{f}_i \hat{\omega}_j^{-1} &= \hat{f}_i \langle \omega_i', \omega_j \rangle^{-1}, \\ [\hat{e}_i, \hat{f}_j] &= \delta_{ij} \frac{\hat{\omega}_i - \hat{\omega}_i'}{r_i - s_i}. \end{aligned}$$

The coefficients  $\langle \omega'_i, \omega_j \rangle^{-1}$  of  $\hat{f}_i$  in the third formula above coincide with those of the formulae related to  $f_i$  in (X2) (here  $X = A, B, C, D$ , resp.). By Lemma 1.2, we find that the coefficients  $\langle \omega'_j, \omega_i \rangle^{-1}$  of  $\hat{e}_i$  in the second formula above do coincide with those of the formulae related to  $e_i$  in (X3). Hence, applying  $\varphi$  gives the desired relations (X2), (X3) and (X4) in  $U$ .

As  $U$  is generated by  $e_i, f_i, \omega_i^{\pm 1}$  and  $\omega'_i{}^{\pm 1}$  ( $1 \leq i \leq n$ ), the mapping  $\varphi$  is surjective. Since  $U$  and  $\mathcal{D}(\mathcal{B}, \mathcal{B}')$  are universal, subject to the same relations on essentially the same generating set,  $\varphi$  provides an isomorphism.  $\square$

**Remarks.** (1) Up to now, we have completely solved the compatibility problem on the defining relations of our two-parameter quantum groups  $U_{r,s}(\mathfrak{g})$  for  $\mathfrak{g} = \mathfrak{so}_{2n+1}, \mathfrak{so}_{2n}$  and  $\mathfrak{sp}_{2n}$ . This is done in two steps: the proof of Theorem 2.5 indicates that the cross relations between  $\mathcal{B}$  and  $\mathcal{B}'$  are half of the relations (X1)—(X4), and the proof of Proposition 2.3 shows the remaining relations, including the remaining half of (X1)—(X4) and the  $(r, s)$ -Serre relations (X5)—(X7).

(2) Let  $U^0 = \mathbb{Q}(r, s)[\omega_1^{\pm 1}, \dots, \omega_n^{\pm 1}, \omega'_1{}^{\pm 1}, \dots, \omega'_n{}^{\pm 1}]$ ,  $U_0 = \mathbb{Q}(r, s)[\omega_1^{\pm 1}, \dots, \omega_n^{\pm 1}]$ , and  $U'_0 = \mathbb{Q}(r, s)[\omega'_1{}^{\pm 1}, \dots, \omega'_n{}^{\pm 1}]$  denote the Laurent polynomial subalgebras of  $U_{r,s}(\mathfrak{g})$ ,  $\mathcal{B}$ , and  $\mathcal{B}'$  respectively. Clearly,  $U^0 = U_0 U'_0 = U'_0 U_0$ . Furthermore, let us denote by  $U_{r,s}(\mathfrak{n})$  (resp.  $U_{r,s}(\mathfrak{n}^-)$ ) the subalgebra of  $\mathcal{B}$  (resp.  $\mathcal{B}'$ ) generated by  $e_i$  (resp.  $f_i$ ) for all  $i \leq n$ . Thus, by definition, we have  $\mathcal{B} = U_{r,s}(\mathfrak{n}) \rtimes U_0$ , and  $\mathcal{B}' = U'_0 \rtimes U_{r,s}(\mathfrak{n}^-)$ , so that the double  $\mathcal{D}(\mathcal{B}, \mathcal{B}') \cong U_{r,s}(\mathfrak{n}) \otimes U^0 \otimes U_{r,s}(\mathfrak{n}^-)$ , as vector spaces.

(3) The above Theorem further implies the existence of the following standard triangular decomposition of  $U_{r,s}(\mathfrak{g})$ , which means that  $U_{r,s}(\mathfrak{g})$  possesses highest weight representation theory in the usual sense. In the corollary below, we point out that the standard triangular decomposition structure of  $U_{r,s}(\mathfrak{g})$  in “two-parameter” is a natural consequence of its Drinfel’d double structure. This yields a direct approach to the triangular decomposition structure of the “one-parameter” quantum groups  $U_q(\mathfrak{g})$  of Drinfel’d-Jimbo type. The two original proofs by Lusztig [16] and Rosso [20] were remarkable but nontrivial compared to ours.

**Corollary 2.6.**  $U_{r,s}(\mathfrak{g}) \cong U_{r,s}(\mathfrak{n}^-) \otimes U^0 \otimes U_{r,s}(\mathfrak{n})$ , as vector spaces. In particular, it induces  $U_q(\mathfrak{g}) \cong U_q(\mathfrak{n}^-) \otimes U_0 \otimes U_q(\mathfrak{n})$ , as vector spaces.

*Proof.* Define  $\langle \cdot, \cdot \rangle^- : \mathcal{B}' \times \mathcal{B} \longrightarrow \mathbb{Q}(r, s)$  by  $\langle b', b \rangle^- := \langle S(b'), b \rangle$  for  $b' \in \mathcal{B}'$ ,  $b \in \mathcal{B}$ , which is the convolution inverse of the skew-dual pairing  $\langle \cdot, \cdot \rangle$  in Proposition 2.3. By definition, it is easily seen that its composition with the flip mapping  $\tau$  yields a new skew-dual pairing  $\langle | \rangle_0 := \langle \cdot, \cdot \rangle^- \circ \tau : \mathcal{B} \times \mathcal{B}' \longrightarrow \mathbb{Q}(r, s)$ , given by  $\langle b | b' \rangle_0 = \langle S(b'), b \rangle$ .

Denote  $\mathcal{D}(\mathcal{B}', \mathcal{B})$  the Drinfel’d double constructed from the skew-dual pairing  $\langle | \rangle_0$ . Then the following mapping  $\theta$  (see [13, 8.2.1(26)]) establishes an isomorphism between Hopf algebras  $\mathcal{D}(\mathcal{B}, \mathcal{B}')$  and  $\mathcal{D}(\mathcal{B}', \mathcal{B})$ :

$$\theta(b \otimes b') = \sum \langle b'_{(1)}, b_{(1)} \rangle b'_{(2)} \otimes b_{(2)} \langle b_{(3)} | b'_{(3)} \rangle_0, \quad b \in \mathcal{B}, \quad b' \in \mathcal{B}'.$$

Hence, the composition  $\theta \circ \varphi^{-1}$  yields the required Hopf algebra isomorphism, where  $\varphi$  is used in the proof of Theorem 2.5.

If one takes  $r = q$ ,  $s = q^{-1}$ , the above triangular decomposition of  $U_{r,s}(\mathfrak{g})$ , modulo some identifications in  $U^0$ , implies the “one-parameter” case.  $\square$

Let  $Q = \mathbb{Z}\Psi$  denote the root lattice and set  $Q^+ = \sum_{i=1}^n \mathbb{Z}_{\geq 0} \alpha_i$ . Then for any  $\zeta = \sum_{i=1}^n \zeta_i \alpha_i \in Q$ , we adopt the notation

$$\omega_\zeta = \omega_1^{\zeta_1} \cdots \omega_n^{\zeta_n}, \quad \omega'_\zeta = (\omega'_1)^{\zeta_1} \cdots (\omega'_n)^{\zeta_n}.$$

We obtain a  $Q$ -graded structure on  $U_{r,s}(\mathfrak{g})$  as another Corollary of Proposition 2.3 and Theorem 2.5. This will be useful for the representation theory discussed later.

**Corollary 2.7.** *For any  $\zeta = \sum_{i=1}^n \zeta_i \alpha_i \in Q$ , the defining relations (X2) and (X3) in  $U_{r,s}(\mathfrak{g})$  take the form:*

$$\begin{aligned} \omega_\zeta e_i \omega_\zeta^{-1} &= \langle \omega'_i, \omega_\zeta \rangle e_i, & \omega_\zeta f_i \omega_\zeta^{-1} &= \langle \omega'_i, \omega_\zeta \rangle^{-1} f_i, \\ \omega'_\zeta e_i \omega'_\zeta^{-1} &= \langle \omega'_\zeta, \omega_i \rangle^{-1} e_i, & \omega'_\zeta f_i \omega'_\zeta^{-1} &= \langle \omega'_\zeta, \omega_i \rangle f_i. \end{aligned}$$

$U_{r,s}(\mathfrak{n}^\pm) = \bigoplus_{\eta \in Q^+} U_{r,s}^{\pm\eta}(\mathfrak{n}^\pm)$  is then  $Q^\pm$ -graded with

$$U_{r,s}^\eta(\mathfrak{n}^\pm) = \left\{ a \in U_{r,s}(\mathfrak{n}^\pm) \mid \omega_\zeta a \omega_\zeta^{-1} = \langle \omega'_\eta, \omega_\zeta \rangle a, \omega'_\zeta a \omega'_\zeta^{-1} = \langle \omega'_\zeta, \omega_\eta \rangle^{-1} a \right\}, \quad (12)$$

for  $\eta \in Q^+ \cup Q^-$ .

Furthermore,  $U = \bigoplus_{\eta \in Q} U_{r,s}^\eta(\mathfrak{g})$  is  $Q$ -graded with

$$\begin{aligned} U_{r,s}^\eta(\mathfrak{g}) &= \left\{ \sum F_\alpha \omega'_\mu \omega_\nu E_\beta \in U \mid \omega_\zeta (F_\alpha \omega'_\mu \omega_\nu E_\beta) \omega_\zeta^{-1} = \langle \omega'_{\beta-\alpha}, \omega_\zeta \rangle F_\alpha \omega'_\mu \omega_\nu E_\beta, \right. \\ &\quad \left. \omega'_\zeta (F_\alpha \omega'_\mu \omega_\nu E_\beta) \omega'_\zeta^{-1} = \langle \omega'_\zeta, \omega_{\beta-\alpha} \rangle^{-1} F_\alpha \omega'_\mu \omega_\nu E_\beta, \text{ with } \beta - \alpha = \eta \right\}, \quad (13) \end{aligned}$$

where  $F_\alpha$  (resp.  $E_\beta$ ) runs over monomials  $f_{i_1} \cdots f_{i_l}$  (resp.  $e_{j_1} \cdots e_{j_m}$ ) such that  $\alpha_{i_1} + \cdots + \alpha_{i_l} = \alpha$  (resp.  $\alpha_{j_1} + \cdots + \alpha_{j_m} = \beta$ ).  $\square$

Let  $\langle | \rangle_0 : \mathcal{B} \times \mathcal{B}' \longrightarrow \mathbb{Q}(r, s)$  denote the skew-dual pairing introduced in the proof of Corollary 2.6. Then we have

**Definition 2.8.** *The bilinear form  $\langle , \rangle_U$  on  $U_{r,s}(\mathfrak{g}) \times U_{r,s}(\mathfrak{g})$  defined by*

$$\begin{aligned} \langle F_\alpha \omega'_\mu \omega_\nu E_\beta, F_\theta \omega'_\sigma \omega_\delta E_\gamma \rangle_U &= \langle S(\omega_\nu E_\beta) | F_\theta \omega'_\sigma \rangle_0 \langle \omega_\delta E_\gamma | S(F_\alpha \omega'_\mu) \rangle_0 \\ &= \langle \omega_\nu | \omega'_\sigma \rangle_0^{-1} \langle \omega_\delta | \omega'_\mu \rangle_0^{-1} \langle S(E_\beta) | F_\theta \rangle_0 \langle E_\gamma | S(F_\alpha) \rangle_0 \\ &= \langle \omega'_\sigma, \omega_\nu \rangle \langle \omega'_\mu, \omega_\delta \rangle \langle F_\theta, E_\beta \rangle \langle S^2(F_\alpha), E_\gamma \rangle \end{aligned} \quad (14)$$

is called the Rosso form of the two-parameter quantum group  $U_{r,s}(\mathfrak{g})$ .

**Proposition 2.9.** *The Rosso form  $\langle , \rangle_U$  on  $U_{r,s}(\mathfrak{g}) \times U_{r,s}(\mathfrak{g})$  is  $\text{ad}_l$ -invariant, that is,*

$$\langle \text{ad}_l(a) b, c \rangle_U = \langle b, \text{ad}_l(S(a)) c \rangle_U, \quad a, b, c \in U_{r,s}(\mathfrak{g}).$$

*Proof.* By Corollary 2.6,  $U_{r,s}(\mathfrak{g}) \cong \mathcal{D}(\mathcal{B}', \mathcal{B})$  with respect to  $\langle | \rangle_0$ . By [13, Proposition 8.12], the Rosso form  $\langle \cdot, \cdot \rangle_U$  on the quantum double  $\mathcal{D}(\mathcal{B}', \mathcal{B})$  is  $\text{ad}_l$ -invariant.  $\square$

### 3. LUSZTIG'S SYMMETRY

We define in this section the Lusztig's symmetries for the two-parameter quantum groups  $U_{r,s}(\mathfrak{g})$  we have defined in Section 1. A remarkable feature of these symmetries in the “two-parameter” cases is their existence between these quantum groups and the so-called associated objects only as  $\mathbb{Q}$ -isomorphisms rather than as  $\mathbb{Q}(r,s)$ -automorphisms as usual in the “one-parameter” cases. When  $\text{rank}(\mathfrak{g}) > 2$ , we find that we can construct the Lusztig's symmetries of  $U_{r,s}(\mathfrak{g})$  into its associated quantum group  $U_{s^{-1},r^{-1}}(\mathfrak{g})$  (as defined below) if and only if  $rs = 1$ . In this case, if we set  $r = q$  and  $s = q^{-1}$ , the Lusztig's symmetries turn out to be  $\mathbb{Q}(q)$ -automorphisms of  $U_{q,q^{-1}}(\mathfrak{g})$ , which in particular induce the usual Lusztig's symmetries defined on quantum groups  $U_q(\mathfrak{g})$  of Drinfel'd-Jimbo type.

**Theorem 3.1.** (i) *When  $\text{rank}(\mathfrak{g}) = 2$ , for  $\mathfrak{g} = \mathfrak{sl}_3$ ,  $\mathfrak{sp}_4$ , or  $\mathfrak{so}_5$ , the Lusztig's symmetries exist between  $U_{r,s}(\mathfrak{g})$  and its associated quantum group  $U_{s^{-1},r^{-1}}(\mathfrak{g})$ .*

(ii) *When  $\text{rank}(\mathfrak{g}) > 2$  for any type of  $\mathfrak{g}$ , the two-parameter quantum group  $U_{r,s}(\mathfrak{g})$  has the Lusztig's symmetries if and only if it is of the form  $U_{q,q^{-1}}(\mathfrak{g})$ , where  $r = q$ ,  $s = q^{-1}$ . In particular, in this case, each Lusztig's symmetry induces a usual Lusztig's symmetry on  $U_q(\mathfrak{g})$ .*

Before giving the proof, we need to make some general preliminary remarks. We first observe that the pairing  $\langle \cdot, \cdot \rangle$  in Proposition 2.3 plays a role in locating the structure constants of  $U_{r,s}(\mathfrak{g})$ . When it is necessary to emphasize this point, we denote the two-parameter quantum groups by  $(U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$ . Now we call  $(U_{s^{-1},r^{-1}}(\mathfrak{g}), \langle | \cdot \rangle)$  the associated quantum group corresponding to  $(U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$ , where the pairing  $\langle \omega'_i | \omega_j \rangle$  is defined via substituting  $(r, s)$  by  $(s^{-1}, r^{-1})$  in the defining formula for  $\langle \omega'_i, \omega_j \rangle$ . Lemma 1.2 guarantees that

$$\langle \omega'_i | \omega_j \rangle = \langle \omega'_j, \omega_i \rangle. \quad (1)$$

In order to define the Lusztig's symmetries, we introduce the notion of divided-power elements: for any nonnegative integer  $k \in \mathbb{N}$ , set

$$\begin{aligned} [k] &= \frac{r^k - s^k}{r - s}, & [k]! &= [1][2] \cdots [k], \\ \langle k \rangle &= \frac{s^{-k} - r^{-k}}{s^{-1} - r^{-1}}, & \langle k \rangle! &= \langle 1 \rangle \langle 2 \rangle \cdots \langle k \rangle, \end{aligned}$$

and for any element  $x \in U_{r,s}(\mathfrak{g})$  (or  $U_{s^{-1},r^{-1}}(\mathfrak{g})$ ), we define two kinds of divided-power elements:

$$x^{[k]} = x^k / [k]!, \quad x^{(k)} = x^k / \langle k \rangle!.$$

**Definition 3.2.** *To every  $i$ ,  $i = 1, \dots, n$ , there corresponds a  $\mathbb{Q}$ -linear mapping  $\mathcal{T}_i : (U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle) \longrightarrow (U_{s^{-1},r^{-1}}(\mathfrak{g}), \langle | \cdot \rangle)$  defined on generators  $\omega_j, \omega'_j, e_j, f_j$  as follow:*

$$\begin{aligned} \mathcal{T}_i(\omega_j) &= \omega_j \omega_i^{-a_{ij}}, & \mathcal{T}_i(\omega'_j) &= \omega'_j \omega_i'^{-a_{ij}}, \\ \mathcal{T}_i(e_i) &= -\omega_i'^{-1} f_i, & \mathcal{T}_i(f_i) &= -(r_i s_i) e_i \omega_i^{-1}; \end{aligned}$$

when  $\mathfrak{g}$  is of type  $A$ ,  $C$ , or  $D$ ,

$$\begin{aligned} \mathcal{T}_i(e_j) &= \sum_{\nu=0}^{-a_{ij}} (-1)^\nu (rs)^{\frac{\nu}{2}(-a_{ij}-\nu)} \langle \omega'_j, \omega_i \rangle^{-\nu} \langle \omega'_i, \omega_i \rangle^{\frac{\nu}{2}(1+a_{ij})} e_i^{(\nu)} e_j e_i^{(-a_{ij}-\nu)}, \quad i \neq j, \\ \mathcal{T}_i(f_j) &= (r_j s_j)^{\delta_{ij}^+} \sum_{\nu=0}^{-a_{ij}} (-1)^\nu (rs)^{\frac{\nu}{2}(-a_{ij}-\nu)} \langle \omega'_i, \omega_j \rangle^\nu \langle \omega'_i, \omega_i \rangle^{-\frac{\nu}{2}(1+a_{ij})} f_i^{(-a_{ij}-\nu)} f_j f_i^{(\nu)}, \quad i \neq j; \end{aligned}$$

and when  $\mathfrak{g}$  is of type  $B$ ,

$$\begin{aligned} \mathcal{T}_i(e_j) &= \sum_{\nu=0}^{-a_{ij}} (-1)^\nu (rs)^{\frac{\nu}{2}(-a_{ij}-\nu)} \langle \omega'_j, \omega_i \rangle^\nu \langle \omega'_i, \omega_i \rangle^{-\frac{\nu}{2}(1+a_{ij})} e_i^{(-a_{ij}-\nu)} e_j e_i^{(\nu)}, \quad i \neq j, \\ \mathcal{T}_i(f_j) &= (r_j s_j)^{\delta_{ij}^-} \sum_{\nu=0}^{-a_{ij}} (-1)^\nu (rs)^{\frac{\nu}{2}(-a_{ij}-\nu)} \langle \omega'_i, \omega_j \rangle^{-\nu} \langle \omega'_i, \omega_i \rangle^{\frac{\nu}{2}(1+a_{ij})} f_i^{(\nu)} f_j f_i^{(-a_{ij}-\nu)}, \quad i \neq j. \end{aligned}$$

Furthermore  $\mathcal{T}_i(r) = s^{-1}$  and  $\mathcal{T}_i(s) = r^{-1}$  so that  $\mathcal{T}_i(x^{[k]}) = (\mathcal{T}_i(x))^{(k)}$ . Here  $(a_{ij})$  is the Cartan matrix of the classical simple Lie algebra  $\mathfrak{g}$ , and for any  $i \neq j$ ,

$$\begin{aligned} \delta_{ij}^+ &= \begin{cases} 2, & \text{if } i < j, \text{ \& } a_{ij} \neq 0, \\ 1, & \text{otherwise.} \end{cases} & \delta_{ij}^- &= \begin{cases} 2, & \text{if } i > j, \text{ \& } a_{ij} \neq 0, \\ 1, & \text{otherwise.} \end{cases} \\ e_i^{(k)} &= e_i^k / \langle k \rangle_i!, & f_i^{(k)} &= f_i^k / \langle k \rangle_i!, & \langle k \rangle_i &= \frac{s_i^{-k} - r_i^{-k}}{s_i^{-1} - r_i^{-1}}. \end{aligned}$$

**Remarks.** (1) The defining formulas of the actions of  $\mathcal{T}_i$  on  $e_j$  and  $f_j$  when  $i \neq j$  (in the cases of types  $A$ ,  $C$ , and  $D$ ) can be interpreted as the right- and the left-adjoint actions of the divided-power operators  $(\text{ad}_r e_i)^{(-a_{ij})}$  and  $(\text{ad}_l f_i)^{(-a_{ij})}$  in the co-opposite Hopf algebra  $U_{r,s}(\mathfrak{g})^{\text{cop}}$ :

$$\mathcal{T}_i(e_j) = (\text{ad}_r e_i)^{(-a_{ij})}(e_j), \quad \mathcal{T}_i(f_j) = (r_j s_j)^{\delta_{ij}^+} (\text{ad}_l f_i)^{(-a_{ij})}(f_j).$$

This phenomenon for the “two-parameter” cases is interesting as it implies that  $U_{r,s}(\mathfrak{g})^{\text{cop}} \cong U_{s^{-1}, r^{-1}}(\mathfrak{g})$  as Hopf algebras. While for the case of type  $B$ , it can be roughly viewed as the dual case of type  $C$ .

According to Definition 3.2, we can also consider the Lusztig’s symmetries  $\mathcal{T}_i$  defined for  $U_{r,s}(\mathfrak{g})$  as  $\mathbb{Q}$ -automorphisms of  $U_{r,s}(\mathfrak{g})$  into itself. However, the images of  $\mathcal{T}_i$  should be contained “locally” (here we mean only in each rank 2 size) in its associated quantum group  $U_{s^{-1}, r^{-1}}(\mathfrak{g})$ . This fact will be clear, in light of the proofs of Lemmas 3.3 — 3.5 below.

(2) When  $r = s^{-1} = q$ , the Lusztig’s symmetries  $\mathcal{T}_i$  are  $\mathbb{Q}(q)$ -automorphisms of  $U_{q,q^{-1}}(\mathfrak{g})$ . This is clear if we extend the action of  $\mathcal{T}_i$  on  $U_{r,s}(\mathfrak{g})$  algebraically, via an identification, we then get the usual Lusztig’s symmetries defined on the quantum groups  $U_q(\mathfrak{g})$  of Drinfel’d-Jimbo type.

In the following, we consider the special cases in rank 2:  $A_2$ ,  $B_2$ , or  $C_2$ .

**Lemma 3.3.** *Assume that  $\mathfrak{g} = \mathfrak{sl}_3$ ,  $\mathfrak{so}_5$ , or  $\mathfrak{sp}_4$ . Then  $\mathcal{T}_i$  ( $i = 1, 2$ ) preserve the defining relations (X1)—(X3) of  $(U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$  into its associated object  $(U_{s^{-1}, r^{-1}}(\mathfrak{g}), \langle \cdot | \cdot \rangle)$ , where  $X = A, B$ , or  $C$ .*

*Proof.* In case  $(A_2)$ : we have

$$\begin{aligned}\langle \omega'_1, \omega_1 \rangle &= rs^{-1} = \langle \omega'_1 | \omega_1 \rangle, & \langle \omega'_1, \omega_2 \rangle &= r^{-1} = \langle \omega'_2 | \omega_1 \rangle, \\ \langle \omega'_2, \omega_1 \rangle &= s = \langle \omega'_1 | \omega_2 \rangle, & \langle \omega'_2, \omega_2 \rangle &= rs^{-1} = \langle \omega'_2 | \omega_2 \rangle.\end{aligned}$$

In case  $(B_2)$ : we have

$$\begin{aligned}\langle \omega'_1, \omega_1 \rangle &= (rs^{-1})^2 = \langle \omega'_1 | \omega_1 \rangle, & \langle \omega'_1, \omega_2 \rangle &= r^{-2} = \langle \omega'_2 | \omega_1 \rangle, \\ \langle \omega'_2, \omega_1 \rangle &= s^2 = \langle \omega'_1 | \omega_2 \rangle, & \langle \omega'_2, \omega_2 \rangle &= rs^{-1} = \langle \omega'_2 | \omega_2 \rangle.\end{aligned}$$

In case  $(C_2)$ : we have

$$\begin{aligned}\langle \omega'_1, \omega_1 \rangle &= rs^{-1} = \langle \omega'_1 | \omega_1 \rangle, & \langle \omega'_1, \omega_2 \rangle &= r^{-2} = \langle \omega'_2 | \omega_1 \rangle, \\ \langle \omega'_2, \omega_1 \rangle &= s^2 = \langle \omega'_1 | \omega_2 \rangle, & \langle \omega'_2, \omega_2 \rangle &= r^2 s^{-2} = \langle \omega'_2 | \omega_2 \rangle.\end{aligned}$$

We need to show that  $\mathcal{T}_1, \mathcal{T}_2$  preserve the defining relations (X1)—(X3). (X1) are automatically satisfied.

To check (X2) & (X3) we first remark that in the rank 2 cases, we have  $\mathcal{T}_k(\langle \omega'_i, \omega_j \rangle) = \langle \mathcal{T}_k(\omega'_i), \mathcal{T}_k(\omega_j) \rangle = \langle \omega'_j, \omega_i \rangle = \langle \omega'_i | \omega_j \rangle$ , for  $i, j, k \in \{1, 2\}$ . This fact ensures that  $\mathcal{T}_k$  ( $k = 1, 2$ ) preserve (X2) and (X3), that is

$$\begin{aligned}\mathcal{T}_k(\omega_j) \mathcal{T}_k(e_i) \mathcal{T}_k(\omega_j)^{-1} &= \langle \omega'_i | \omega_j \rangle \mathcal{T}_k(e_i), & \mathcal{T}_k(\omega_j) \mathcal{T}_k(f_i) \mathcal{T}_k(\omega_j)^{-1} &= \langle \omega'_i | \omega_j \rangle^{-1} \mathcal{T}_k(f_i), \\ \mathcal{T}_k(\omega'_j) \mathcal{T}_k(e_i) \mathcal{T}_k(\omega'_j)^{-1} &= \langle \omega'_j | \omega_i \rangle^{-1} \mathcal{T}_k(e_i), & \mathcal{T}_k(\omega'_j) \mathcal{T}_k(f_i) \mathcal{T}_k(\omega'_j)^{-1} &= \langle \omega'_j | \omega_i \rangle \mathcal{T}_k(f_i).\end{aligned}$$

All identities follow from the first one.  $\square$

**Lemma 3.4.** *With the same assumption as in Lemma 3.3,  $\mathcal{T}_i$  ( $i = 1, 2$ ) preserves the defining relations (X4) of  $(U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$  into its associated object  $(U_{s^{-1}, r^{-1}}(\mathfrak{g}), \langle \cdot | \cdot \rangle)$ , for  $X = A, B, C$ .*

*Proof.* For  $i = 1, 2$ , we have

$$\begin{aligned}[\mathcal{T}_i(e_i), \mathcal{T}_i(f_i)] &= (r_i s_i) \omega_i'^{-1} (f_i e_i - \langle \omega'_i, \omega_i \rangle^{-1} \langle \omega'_i, \omega_i \rangle e_i f_i) \omega_i'^{-1} \\ &= \frac{\omega_i^{-1} - \omega_i'^{-1}}{s_i^{-1} - r_i^{-1}} = \frac{\mathcal{T}_i(\omega_i) - \mathcal{T}_i(\omega'_i)}{s_i^{-1} - r_i^{-1}} \\ &= \mathcal{T}_i([e_i, f_i]) \in U_{s^{-1}, r^{-1}}(\mathfrak{g}).\end{aligned}$$

When  $a_{ij} = -1$ , we have

$$\begin{aligned}
[\mathcal{T}_i(e_j), \mathcal{T}_i(f_j)] &= (r_j s_j)^{\delta_{ij}^+} [e_j e_i - \langle \omega'_j, \omega_i \rangle^{-1} e_i e_j, f_i f_j - \langle \omega'_i, \omega_j \rangle f_j f_i] \\
&= (r_j s_j)^{\delta_{ij}^+} \{ f_i [e_j, f_j] e_i + e_j [e_i, f_i] f_j - \langle \omega'_j, \omega_i \rangle^{-1} ([e_i, f_i] f_j e_j + e_i f_i [e_j, f_j]) \\
&\quad - \langle \omega'_i, \omega_j \rangle ([e_j, f_j] f_i e_i + e_j f_j [e_i, f_i]) \\
&\quad + \langle \omega'_j, \omega_i \rangle^{-1} \langle \omega'_i, \omega_j \rangle (e_i [e_j, f_j] f_i + f_j [e_i, f_i] e_j) \} \\
&= \frac{(r_j s_j)^{\delta_{ij}^+}}{r_j - s_j} \frac{1}{r_i - s_i} \{ (\omega_j - \omega'_j) [\omega_i \langle \omega'_j, \omega_i \rangle^{-1} - \omega'_i \langle \omega'_i, \omega_j \rangle \\
&\quad - (\langle \omega'_j, \omega_i \rangle^{-1} + \langle \omega'_i, \omega_j \rangle) (\omega_i - \omega'_i)] \\
&\quad + \langle \omega'_j, \omega_i \rangle^{-1} \langle \omega'_i, \omega_j \rangle (\omega_j \langle \omega'_i, \omega_j \rangle^{-1} - \omega'_j \langle \omega'_j, \omega_i \rangle) (\omega_i - \omega'_i) \} \\
&= \frac{(r_j s_j)^{\delta_{ij}^+} (\langle \omega'_j, \omega_i \rangle^{-1} - \langle \omega'_i, \omega_j \rangle)}{(r_j - s_j)(r_i - s_i)} (\omega_i \omega_j - \omega'_i \omega'_j) \\
&= \begin{cases} \frac{\mathcal{T}_i(\omega_j) - \mathcal{T}_i(\omega'_j)}{s^{-1} - r^{-1}}, & \text{for } (i, j) = (1, 2), \text{ i.e., in type } A_2 \\ \frac{\mathcal{T}_i(\omega_j) - \mathcal{T}_i(\omega'_j)}{s^{-1} - r^{-1}}, & \text{for } (i, j) = (2, 1), \text{ i.e., in types } A_2, C_2 \end{cases} \\
&= \mathcal{T}_i([e_j, f_j]).
\end{aligned}$$

When  $a_{12} = -2$  in type  $C_2$ , we need to prove

$$[\mathcal{T}_1(e_2), \mathcal{T}_1(f_2)] = \frac{\omega_2 \omega_1^2 - \omega'_2 \omega_1'^2}{s^{-2} - r^{-2}} = \frac{\mathcal{T}_1(\omega_2) - \mathcal{T}_1(\omega'_2)}{s_2^{-1} - r_2^{-1}} = \mathcal{T}_1([e_2, f_2]) \in U_{s^{-1}, r^{-1}}(\mathfrak{g}).$$

To this end, let us denote

$$\begin{aligned}
E_{12} &= e_{\alpha_1 + \alpha_2} = \text{ad}_l e_1(e_2) = e_1 e_2 - s^2 e_2 e_1, \\
F_{12} &= f_{\alpha_1 + \alpha_2} = \text{ad}_r f_1(f_2) = f_2 f_1 - r^2 f_1 f_2, \\
E_{112} &= e_{2\alpha_1 + \alpha_2} = \text{ad}_l e_1^2(e_2) = e_1 E_{12} - r s E_{12} e_1, \\
F_{112} &= f_{2\alpha_1 + \alpha_2} = \text{ad}_r f_1^2(f_2) = F_{12} f_1 - r s f_1 F_{12}.
\end{aligned}$$

A direct calculation shows

$$\begin{aligned}
[e_1, F_{12}] &= -(r + s) \omega_1 f_2, & [e_2, F_{12}] &= f_1 \omega'_2, \\
[E_{12}, f_1] &= -(r + s) e_2 \omega'_1, & [E_{12}, f_2] &= \omega_2 e_1, \\
[E_{12}, F_{12}] &= \frac{\omega_1 \omega_2 - \omega'_1 \omega'_2}{r - s}.
\end{aligned}$$

Using the Leibniz rule and the above results, we can get

$$\begin{aligned}
[E_{112}, F_{112}] &= [e_1 E_{12} - r s E_{12} e_1, F_{12} f_1 - r s f_1 F_{12}] \\
&= [e_1, F_{12}] f_1 E_{12} + e_1 [E_{12}, F_{12}] f_1 + F_{12} [e_1, f_1] E_{12} + e_1 F_{12} [E_{12}, f_1] \\
&\quad - (r s) ([E_{12}, F_{12}] f_1 e_1 + E_{12} [e_1, F_{12}] f_1 + F_{12} [E_{12}, f_1] e_1 + E_{12} F_{12} [e_1, f_1]) \\
&\quad - (r s) ([e_1, f_1] F_{12} E_{12} + f_1 [e_1, F_{12}] E_{12} + e_1 [E_{12}, f_1] F_{12} + e_1 f_1 [E_{12}, F_{12}]) \\
&\quad + (r s)^2 (E_{12} [e_1, f_1] F_{12} + E_{12} f_1 [e_1, F_{12}] + [E_{12}, f_1] F_{12} e_1 + f_1 [E_{12}, F_{12}] e_1) \\
&= \frac{(r + s)^2}{r^2 - s^2} (\omega_1^2 \omega_2 - \omega_1'^2 \omega_2').
\end{aligned}$$

Observe that  $\mathcal{T}_1(e_2) = \frac{1}{\langle 2 \rangle_1} E_{112}$ ,  $\mathcal{T}_1(f_2) = \frac{1}{\langle 2 \rangle_1} F_{112}$ , where  $\langle 2 \rangle_1 = (r+s)(rs)^{-1}$ . We then arrive at the required equality above.

A similar argument is used to prove the result for type  $B_2$ .  $\square$

**Lemma 3.5.** *With the same assumption as in Lemma 3.3,  $\mathcal{T}_i$  ( $i = 1, 2$ ) preserves the  $(r, s)$ -Serre relations (X5), (X6) and (X7) of  $(U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$  into its associated object  $(U_{s^{-1}, r^{-1}}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$ , for  $X = A, B, C$ .*

*Proof.* We do not need to consider the relation (X5) as it doesn't appear for the rank 2 cases.

For  $A_2$ : Consider the degree 2  $(r, s)$ -Serre relation

$$e_1^2 e_2 - (r+s)e_1 e_2 e_1 + (rs)e_2 e_1^2 = 0. \quad (2)$$

Note that

$$\begin{aligned} \mathcal{T}_1(e_2)\mathcal{T}_1(e_1) &= s\mathcal{T}_1(e_1)\mathcal{T}_1(e_2) - r^{-1}e_2, & e_2\mathcal{T}_1(e_1) &= r\mathcal{T}_1(e_1)e_2; \\ \mathcal{T}_2(e_1)\mathcal{T}_2(e_2) &= r^{-1}\mathcal{T}_2(e_2)\mathcal{T}_2(e_1) - r^{-1}e_1, & e_1\mathcal{T}_2(e_1) &= s\mathcal{T}_2(e_1)e_1. \end{aligned}$$

Acting with  $\mathcal{T}_i$  ( $i = 1, 2$ ) algebraically on the left-hand side of (2), we can get

$$\begin{aligned} \mathcal{T}_1(e_1)^2 \mathcal{T}_1(e_2) - (r^{-1} + s^{-1})\mathcal{T}_1(e_1)\mathcal{T}_1(e_2)\mathcal{T}_1(e_1) + (r^{-1}s^{-1})\mathcal{T}_1(e_2)\mathcal{T}_1(e_1)^2 &= 0, \\ \mathcal{T}_2(e_1)^2 \mathcal{T}_2(e_2) - (r^{-1} + s^{-1})\mathcal{T}_2(e_1)\mathcal{T}_2(e_2)\mathcal{T}_2(e_1) + (r^{-1}s^{-1})\mathcal{T}_2(e_2)\mathcal{T}_2(e_1)^2 &= 0, \end{aligned}$$

that is,  $\mathcal{T}_i$  ( $i = 1, 2$ ) preserves (2) for  $A_2$ .

Now consider another degree 2  $(r, s)$ -Serre relation

$$e_2^2 e_1 - (r_2^{-1} + s_2^{-1})e_2 e_1 e_2 + (r_2^{-1}s_2^{-1})e_1 e_2^2 = 0, \quad (3)$$

which holds for both  $A_2$  and  $C_2$ . Note that

$$\mathcal{T}_2(e_1)\mathcal{T}_2(e_2) = r_2^{-1}\mathcal{T}_2(e_2)\mathcal{T}_2(e_1) - r_2^{-1}e_1, \quad \mathcal{T}_2(e_2)e_1 = s_2 e_1 \mathcal{T}_2(e_2).$$

Acting with  $\mathcal{T}_2$  algebraically on the left-hand side of (2), we can easily get

$$\mathcal{T}_2(e_2)^2 \mathcal{T}_2(e_1) - (r_2 + s_2)\mathcal{T}_2(e_2)\mathcal{T}_2(e_1)\mathcal{T}_2(e_2) + (r_2 s_2)\mathcal{T}_2(e_1)\mathcal{T}_2(e_2)^2 = 0,$$

that is,  $\mathcal{T}_2$  preserves (3) in both cases  $A_2$  and  $C_2$ .

In the case of  $A_2$ , let  $\mathcal{T}_1$  act algebraically on the left-hand side of (3). Because

$$\mathcal{T}_1(e_2)\mathcal{T}_1(e_1) = s\mathcal{T}_1(e_1)\mathcal{T}_1(e_2) - r^{-1}e_2, \quad e_2\mathcal{T}_1(e_2) = r^{-1}\mathcal{T}_1(e_2)e_2,$$

we can easily get that

$$\mathcal{T}_1(e_2)^2 \mathcal{T}_1(e_1) - (r+s)\mathcal{T}_1(e_2)\mathcal{T}_1(e_1)\mathcal{T}_1(e_2) + (rs)\mathcal{T}_1(e_1)\mathcal{T}_1(e_2)^2 = 0.$$

That is,  $\mathcal{T}_1$  preserves (3) for  $A_2$ , as well.

For  $C_2$ : Note that  $[e_1^{(2)}, f_1] = \frac{s\omega_1 - r\omega'_1}{r-s}e_1$ . So by direct calculation, we get

$$\begin{aligned}\mathcal{T}_1(e_2)\mathcal{T}_1(e_1) &= s_2\mathcal{T}_1(e_1)\mathcal{T}_1(e_2) + (rs)^{-1}\mathcal{T}'_1(e_2), \\ \mathcal{T}_1(e_2)\mathcal{T}'_1(e_2) &= r^2\mathcal{T}'_1(e_2)\mathcal{T}_1(e_2), \quad (\text{by Lemma 3.6})\end{aligned}\tag{4}$$

where  $\mathcal{T}_1(e_2) = e_2e_1^{(2)} - s^{-1}e_1e_2e_1 + r^{-1}s^{-3}e_1^{(2)}e_2$ ,  $\mathcal{T}'_1(e_2) = (\text{ad}_l e_1)(e_2) = e_1e_2 - s^2e_2e_1$ .

Let  $\mathcal{T}_1$  act algebraically on the left-hand side of (3), we get

$$\mathcal{T}_1(e_2)^2\mathcal{T}_1(e_1) - (r_2 + s_2)\mathcal{T}_1(e_2)\mathcal{T}_1(e_1)\mathcal{T}_1(e_2) + (r_2s_2)\mathcal{T}_1(e_1)\mathcal{T}_1(e_2)^2 = 0.$$

That is,  $\mathcal{T}_1$  preserves (3) for  $C_2$ .

Now consider the degree 3  $(r, s)$ -Serre relation

$$e_1^3e_2 - (r^2 + rs + s^2)e_1^2e_2e_1 + (rs)(r^2 + rs + s^2)e_1e_2e_1^2 - (rs)^3e_2e_1^3 = 0.\tag{5}$$

Observing that

$$\begin{aligned}e_2\mathcal{T}_1(e_1) &= r^2\mathcal{T}_1(e_1)e_2, \\ \mathcal{T}'_1(e_2)\mathcal{T}_1(e_1) &= (rs)\mathcal{T}_1(e_1)\mathcal{T}'_1(e_2) + r^{-1}s(r+s)e_2,\end{aligned}$$

and using (4), it is easy to check that  $\mathcal{T}_1$  preserves (5).

On the other hand, note that

$$\begin{aligned}\mathcal{T}_2(e_1)\mathcal{T}_2(e_2) &= r^{-2}\mathcal{T}_2(e_2)\mathcal{T}_2(e_1) - r^{-2}e_1, \\ e_1\mathcal{T}_2(e_1)^2 - s(r+s)\mathcal{T}_2(e_1)e_1\mathcal{T}_2(e_1) + rs^3\mathcal{T}_2(e_1)^2e_1 &= 0, \quad (\text{by Lemma 3.6})\end{aligned}$$

which ensure  $\mathcal{T}_2$  preserves (5).

Dually, we can verify  $\mathcal{T}_i$  ( $i = 1, 2$ ) preserves the  $(r, s)$ -Serre relations (X7) for  $X = A_2, C_2$ .

We can prove the result for type  $B_2$  in a similar way.  $\square$

**Lemma 3.6.** *For type  $C_2$ , the following identities hold.*

$$\mathcal{T}_1(e_2)\mathcal{T}'_1(e_2) = r^2\mathcal{T}'_1(e_2)\mathcal{T}_1(e_2),\tag{6}$$

$$e_1\mathcal{T}_2(e_1)^2 - s(r+s)\mathcal{T}_2(e_1)e_1\mathcal{T}_2(e_1) + rs^3\mathcal{T}_2(e_1)^2e_1 = 0,\tag{7}$$

where  $\mathcal{T}_1(e_2) = e_2e_1^{(2)} - s^{-1}e_1e_2e_1 + r^{-1}s^{-3}e_1^{(2)}e_2$ ,  $\mathcal{T}'_1(e_2) = (\text{ad}_l e_1)(e_2) = e_1e_2 - s^2e_2e_1$ , and  $\mathcal{T}_2(e_1) = e_1e_2 - r^2e_2e_1$ .

*Proof.* Writing  $\Delta = r^2 + rs + s^2$ , and using (C6), we get

$$\begin{aligned}(rs^3e_2e_1^2 - s(r+s)e_1e_2e_1 + e_1^2e_2)(e_1e_2 - s^2e_2e_1) \\ - r^2(e_1e_2 - s^2e_2e_1)(rs^3e_2e_1^2 - s(r+s)e_1e_2e_1 + e_1^2e_2)\end{aligned}$$

$$\begin{aligned}
&= rs^3 e_2 e_1^3 e_2 - rs^5 e_2 e_1^2 e_2 e_1 - s(r+s) e_1 e_2 e_1^2 e_2 \\
&\quad + s^3(r+s) e_1 e_2 e_1 e_2 e_1 + e_1^2 e_2 e_1 e_2 - s^2 e_1^2 e_2^2 e_1 \\
&\quad - [(rs)^3 e_1 e_2^2 e_1^2 - r^2 s(r+s) e_1 e_2 e_1 e_2 e_1 + r^2 e_1 e_2 e_1^2 e_2 \\
&\quad - r^3 s^5 e_2 e_1 e_2 e_1^2 + r^2 s^3(r+s) e_2 e_1^2 e_2 e_1 - (rs)^2 e_2 e_1^3 e_2] \\
&= rs^2(r+s) e_2 e_1^3 e_2 - rs^3 \Delta e_2(e_1^2 e_2 e_1) - \Delta(e_1 e_2 e_1^2) e_2 + s(r+s)(r^2+s^2) e_1 e_2 e_1 e_2 e_1 \\
&\quad + e_1^2 e_2 e_1 e_2 - s^2 e_1^2 e_2^2 e_1 + r^3 s^5 e_2 e_1 e_2 e_1^2 - (rs)^3 e_1 e_2^2 e_1^2 \\
&= rs^2(r+s) e_2 e_1^3 e_2 - rs^3 e_2 e_1^3 e_2 - r^2 s^4 \Delta e_2 e_1 e_2 e_1^2 + r^4 s^6 e_2^2 e_1^3 \\
&\quad - (rs)^{-1} \Delta e_1^2 e_2 e_1 e_2 + (rs)^{-1} e_1^3 e_2^2 - (rs)^2 e_2 e_1^3 e_2 + s(r+s)(r^2+s^2) e_1 e_2 e_1 e_2 e_1 \\
&\quad + e_1^2 e_2 e_1 e_2 - s^2 e_1^2 e_2^2 e_1 + r^3 s^5 e_2 e_1 e_2 e_1^2 - (rs)^3 e_1 e_2^2 e_1^2 \\
&= -r^2 s^4 (r^2+s^2) (e_2 e_1 e_2) e_1^2 + r^4 s^6 e_2^2 e_1^3 - (rs)^{-1} (r^2+s^2) e_1^2 (e_2 e_1 e_2) \\
&\quad + (rs)^{-1} e_1^3 e_2^2 + s(r+s)(r^2+s^2) e_1 (e_2 e_1 e_2) e_1 - s^2 e_1^2 e_2^2 e_1 - (rs)^3 e_1 e_2^2 e_1^2 \\
&= -r^2 s^4 [(rs)^2 e_2^2 e_1^3 + e_1 e_2^2 e_1^2] + r^4 s^6 e_2^2 e_1^3 - (rs)^{-1} [(rs)^2 e_1^2 e_2^2 e_1 + e_1^3 e_2^2] \\
&\quad + (rs)^{-1} e_1^3 e_2^2 + s(r+s) [(rs)^2 e_1 e_2^2 e_1^2 + e_1^2 e_2^2 e_1] - s^2 e_1^2 e_2^2 e_1 - (rs)^3 e_1 e_2^2 e_1^2 \\
&= 0.
\end{aligned}$$

Thus, we get the identity (6)

$$\mathcal{T}_1(e_2) \mathcal{T}'_1(e_2) = r^2 \mathcal{T}'_1(e_2) \mathcal{T}_1(e_2).$$

To check the identity (7), we have

$$\begin{aligned}
e_1 \mathcal{T}_2(e_1)^2 &= e_1^2 e_2 e_1 e_2 - r^2 e_1^2 e_2^2 e_1 - r^2 e_1 e_2 e_1^2 e_2 + r^4 e_1 e_2 e_1 e_2 e_1, \\
-s(r+s) \mathcal{T}_2(e_1) e_1 \mathcal{T}_2(e_1) &= -s(r+s) e_1 e_2 e_1^2 e_2 + r^2 s(r+s) e_1 e_2 e_1 e_2 e_1 \\
&\quad + r^2 s(r+s) e_2 e_1^3 e_2 - r^4 s(r+s) e_2 e_1^2 e_2 e_1, \\
rs^3 \mathcal{T}_2(e_1)^2 e_1 &= rs^3 e_1 e_2 e_1 e_2 e_1 - (rs)^3 e_1 e_2^2 e_1^2 - (rs)^3 e_2 e_1^2 e_2 e_1 + r^5 s^3 e_2 e_1 e_2 e_1^2.
\end{aligned}$$

So we obtain

$$\begin{aligned}
\text{LHS of (7)} &= e_1^2 e_2 e_1 e_2 - r^2 e_1 (e_1 e_2^2) e_1 - \Delta(e_1 e_2 e_1^2) e_2 + r(r+s)(r^2+s^2) e_1 e_2 e_1 e_2 e_1 \\
&\quad + r^2 s(r+s) e_2 e_1^3 e_2 - (rs)^3 e_1 e_2^2 e_1^2 + r^5 s^3 e_2 e_1 e_2 e_1^2 - r^3 s \Delta e_2(e_1^2 e_2 e_1) \\
&= e_1^2 e_2 e_1 e_2 - r^2 [(r^2+s^2) e_1 e_2 e_1 e_2 e_1 - (rs)^2 e_1 e_2^2 e_1^2] \\
&\quad - [(rs)^2 e_2 e_1^3 e_2 + (rs)^{-1} \Delta e_1^2 e_2 e_1 e_2 - (rs)^{-1} e_1^3 e_2^2] \\
&\quad + r(r+s)(r^2+s^2) e_1 e_2 e_1 e_2 e_1 + r^2 s(r+s) e_2 e_1^3 e_2 - (rs)^3 e_1 e_2^2 e_1^2 \\
&\quad + r^5 s^3 e_2 e_1 e_2 e_1^2 - r^3 s [(rs) \Delta e_2 e_1 e_2 e_1^2 + e_2 e_1^3 e_2 - (rs)^3 e_2^2 e_1^3] \\
&= -(rs)^{-1} (r^2+s^2) e_1^2 (e_2 e_1 e_2) + (rs) (r^2+s^2) e_1 (e_2 e_1 e_2) e_1 + r^3 s^2 (r-s) e_1 e_2^2 e_1^2 \\
&\quad + (rs)^{-1} e_1^3 e_2^2 - r^4 s^2 (r^2+s^2) (e_2 e_1 e_2) e_1^2 + r^6 s^4 e_2^2 e_1^3 \\
&= -(rs)^{-1} [(rs)^2 e_1^2 e_2^2 e_1 + e_1^3 e_2^2] + (rs) [(rs)^2 e_1 e_2^2 e_1^2 + e_1^2 e_2^2 e_1] \\
&\quad + r^3 s^2 (r-s) e_1 e_2^2 e_1^2 + (rs)^{-1} e_1^3 e_2^2 - r^4 s^2 [(rs)^2 e_2^2 e_1^3 + e_1 e_2^2 e_1^2] + r^6 s^4 e_2^2 e_1^3 \\
&= 0.
\end{aligned}$$

This complete the proof.  $\square$

Now let us consider the rank 3 cases.

**Lemma 3.7.** *For the rank 3 cases of types  $A_3$ ,  $B_3$  and  $C_3$ , the Lusztig's symmetries  $\mathcal{T}_i$  preserve the defining relations (X2) or (X3) of  $(U_{r,s}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$  into its associated object  $(U_{s^{-1}, r^{-1}}(\mathfrak{g}), \langle \cdot, \cdot \rangle)$ , for  $X = A, B, C$  if and only if  $rs = 1$ .*

*Proof.* When  $rs = 1$ , we set  $r = q$ ,  $s = q^{-1}$ . It is then obvious that  $U_{q,q^{-1}}(\mathfrak{g})$  has Lusztig's symmetries for any type and any rank of  $\mathfrak{g}$ . Indeed, the quantum group  $U_q(\mathfrak{g})$  of Drinfel'd-Jimbo type is a quotient of  $U_{q,q^{-1}}(\mathfrak{g})$  by the ideal  $(\omega_i^{-1} - \omega'_i, i = 1, \dots, n)$  (see [15]). Consequently, each  $\mathcal{T}_i$  automatically preserves the defining relations (X2) and (X3).

Conversely, assume that each  $\mathcal{T}_i$  preserves relations (X2) or (X3). In type  $A_3$  or  $C_3$  case, we consider the situation where  $\mathcal{T}_3$  preserves  $\omega_1 e_2 \omega_1^{-1} = s e_2$ . This leads to  $\mathcal{T}_3(\omega_1) \mathcal{T}_3(e_2) \mathcal{T}_3(\omega_1^{-1}) = \omega_1(e_2 e_3 - r e_3 e_2) \omega_1^{-1} = s \mathcal{T}_3(e_2) = \mathcal{T}_3(\omega_1 e_2 \omega_1^{-1}) = \mathcal{T}_3(s e_2) = r^{-1} \mathcal{T}_3(e_2)$  and this implies  $rs = 1$ . Similarly, in  $B_3$  case, we can get the same condition provided that we observe that  $\mathcal{T}_1$  preserves the relation  $\omega_3 e_2 \omega_3^{-1} = r^{-1} e_2$ .  $\square$

*Proof of Theorem 3.1.* When  $\text{rank}(\mathfrak{g}) = 2$ , Lemmas 3.3, 3.4 & 3.5 indicate that for arbitrary parameters  $r, s$  with  $r^2 \neq s^2$ ,  $U_{r,s}(\mathfrak{sl}_3)$ ,  $U_{r,s}(\mathfrak{sp}_4)$  and  $U_{r,s}(\mathfrak{so}_5)$  possess the Lusztig symmetries into their respective associated quantum groups. When  $\text{rank}(\mathfrak{g}) > 2$ , this  $\mathfrak{g}$  contains one of rank 3 Lie subalgebras  $\mathfrak{sl}_4$ ,  $\mathfrak{so}_7$  and  $\mathfrak{sp}_6$ . Lemma 3.7 gives the required assertion.  $\square$

#### 4. APPENDIX: SOME CALCULATIONS

**4.1 Calculations in the proof of Proposition 2.3** The relevant terms of  $\Delta(X)$  for  $X = f_1^3 f_2$  in (6C) are as follows:

$$\begin{aligned}
& f_1 \omega_1'^2 \omega_2' \otimes f_1 \omega_1' \omega_2' \otimes f_1 \omega_2' \otimes f_2 + f_1 \omega_1'^2 \omega_2' \otimes \omega_1' f_1 \omega_2' \otimes f_1 \omega_2' \otimes f_2 \\
& \quad + \omega_1' f_1 \omega_1' \omega_2' \otimes f_1 \omega_1' \omega_2' \otimes f_1 \omega_2' \otimes f_2 + \omega_1' f_1 \omega_1' \omega_2' \otimes \omega_1' f_1 \omega_2' \otimes f_1 \omega_2' \otimes f_2 \\
& \quad + \omega_1'^2 f_1 \omega_2' \otimes f_1 \omega_1' \omega_2' \otimes f_1 \omega_2' \otimes f_2 + \omega_1'^2 f_1 \omega_2' \otimes \omega_1' f_1 \omega_2' \otimes f_1 \omega_2' \otimes f_2 \\
& + f_1 \omega_1'^2 \omega_2' \otimes f_1 \omega_1' \omega_2' \otimes \omega_1' f_2 \otimes f_1 + f_1 \omega_1'^2 \omega_2' \otimes \omega_1' f_1 \omega_2' \otimes \omega_1' f_2 \otimes f_1 \\
& \quad + \omega_1' f_1 \omega_1' \omega_2' \otimes f_1 \omega_1' \omega_2' \otimes \omega_1' f_2 \otimes f_1 + \omega_1' f_1 \omega_1' \omega_2' \otimes \omega_1' f_1 \omega_2' \otimes \omega_1' f_2 \otimes f_1 \\
& \quad + \omega_1'^2 f_1 \omega_2' \otimes f_1 \omega_1' \omega_2' \otimes \omega_1' f_2 \otimes f_1 + \omega_1'^2 f_1 \omega_2' \otimes \omega_1' f_1 \omega_2' \otimes \omega_1' f_2 \otimes f_1 \\
& + f_1 \omega_1'^2 \omega_2' \otimes \omega_1'^2 f_2 \otimes f_1 \omega_1' \otimes f_1 + f_1 \omega_1'^2 \omega_2' \otimes \omega_1'^2 f_2 \otimes \omega_1' f_1 \otimes f_1 \\
& \quad + \omega_1' f_1 \omega_1' \omega_2' \otimes \omega_1'^2 f_2 \otimes f_1 \omega_1' \otimes f_1 + \omega_1' f_1 \omega_1' \omega_2' \otimes \omega_1'^2 f_2 \otimes \omega_1' f_1 \otimes f_1 \\
& \quad + \omega_1'^2 f_1 \omega_2' \otimes \omega_1'^2 f_2 \otimes f_1 \omega_1' \otimes f_1 + \omega_1'^2 f_1 \omega_2' \otimes \omega_1'^2 f_2 \otimes \omega_1' f_1 \otimes f_1 \\
& + \omega_1'^3 f_2 \otimes f_1 \omega_1'^2 \otimes f_1 \omega_1' \otimes f_1 + \omega_1'^3 f_2 \otimes f_1 \omega_1'^2 \otimes \omega_1' f_1 \otimes f_1 \\
& \quad + \omega_1'^3 f_2 \otimes \omega_1' f_1 \omega_1' \otimes f_1 \omega_1' \otimes f_1 + \omega_1'^3 f_2 \otimes \omega_1' f_1 \omega_1' \otimes \omega_1' f_1 \otimes f_1 \\
& \quad + \omega_1'^3 f_2 \otimes \omega_1'^2 f_1 \otimes f_1 \omega_1' \otimes f_1 + \omega_1'^3 f_2 \otimes \omega_1'^2 f_1 \otimes \omega_1' f_1 \otimes f_1. \tag{4.1}
\end{aligned}$$

The relevant terms of  $\Delta(X)$  for  $X = f_2 f_1^3$  in (6C) are as follows:

$$\begin{aligned}
& \omega'_1 \omega'_2 f_1 \omega'_1 \otimes \omega'_1 \omega'_2 f_1 \otimes \omega'_2 f_1 \otimes f_2 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_1 \omega'_2 f_1 \otimes \omega'_2 f_1 \otimes f_2 \\
& + \omega'_1 \omega'_2 f_1 \omega'_1 \otimes \omega'_2 f_1 \omega'_1 \otimes \omega'_2 f_1 \otimes f_2 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_2 f_1 \omega'_1 \otimes \omega'_2 f_1 \otimes f_2 \\
& + \omega'_2 f_1 \omega'_1{}^2 \otimes \omega'_2 f_1 \omega'_1 \otimes \omega'_2 f_1 \otimes f_2 + \omega'_2 f_1 \omega'_1{}^2 \otimes \omega'_1 \omega'_2 f_1 \otimes \omega'_2 f_1 \otimes f_2 \\
& + \omega'_1 \omega'_2 f_1 \omega'_1 \otimes \omega'_1 \omega'_2 f_1 \otimes f_2 \omega'_1 \otimes f_1 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_1 \omega'_2 f_1 \otimes f_2 \omega'_1 \otimes f_1 \\
& + \omega'_1 \omega'_2 f_1 \omega'_1 \otimes \omega'_2 f_1 \omega'_1 \otimes f_2 \omega'_1 \otimes f_1 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_2 f_1 \omega'_1 \otimes f_2 \omega'_1 \otimes f_1 \\
& + \omega'_2 f_1 \omega'_1{}^2 \otimes \omega'_2 f_1 \omega'_1 \otimes f_2 \omega'_1 \otimes f_1 + \omega'_2 f_1 \omega'_1{}^2 \otimes \omega'_1 \omega'_2 f_1 \otimes f_2 \omega'_1 \otimes f_1 \\
& + \omega'_2 f_1 \omega'_1{}^2 \otimes f_2 \omega'_1{}^2 \otimes f_1 \omega'_1 \otimes f_1 + \omega'_2 f_1 \omega'_1{}^2 \otimes f_2 \omega'_1{}^2 \otimes \omega'_1 f_1 \otimes f_1 \\
& + \omega'_1 \omega'_2 f_1 \omega'_1 \otimes f_2 \omega'_1{}^2 \otimes f_1 \omega'_1 \otimes f_1 + \omega'_1{}^2 \omega'_2 f_1 \otimes f_2 \omega'_1{}^2 \otimes f_1 \omega'_1 \otimes f_1 \\
& + \omega'_1 \omega'_2 f_1 \omega'_1 \otimes f_2 \omega'_1{}^2 \otimes \omega'_1 f_1 \otimes f_1 + \omega'_1{}^2 \omega'_2 f_1 \otimes f_2 \omega'_1{}^2 \otimes \omega'_1 f_1 \otimes f_1 \\
& + f_2 \omega'_1{}^3 \otimes f_1 \omega'_1{}^2 \otimes f_1 \omega'_1 \otimes f_1 + f_2 \omega'_1{}^3 \otimes f_1 \omega'_1{}^2 \otimes \omega'_1 f_1 \otimes f_1 \\
& + f_2 \omega'_1{}^3 \otimes \omega'_1 f_1 \omega'_1 \otimes f_1 \omega'_1 \otimes f_1 + f_2 \omega'_1{}^3 \otimes \omega'_1{}^2 f_1 \otimes f_1 \omega'_1 \otimes f_1 \\
& + f_2 \omega'_1{}^3 \otimes \omega'_1 f_1 \omega'_1 \otimes \omega'_1 f_1 \otimes f_1 + f_2 \omega'_1{}^3 \otimes \omega'_1{}^2 f_1 \otimes \omega'_1 f_1 \otimes f_1. \tag{4.2}
\end{aligned}$$

The relevant terms of  $\Delta(X)$  for  $X = f_1^2 f_2 f_1$  in (6C) are as follows:

$$\begin{aligned}
& f_1 \omega'_1{}^2 \omega'_2 \otimes f_1 \omega'_1 \omega'_2 \otimes \omega'_2 f_1 \otimes f_2 + \omega'_1 f_1 \omega'_1 \omega'_2 \otimes f_1 \omega'_1 \omega'_2 \otimes \omega'_2 f_1 \otimes f_2 \\
& + f_1 \omega'_1{}^2 \omega'_2 \otimes \omega'_1 \omega'_2 f_1 \otimes f_1 \omega'_2 \otimes f_2 + \omega'_1 f_1 \omega'_1 \omega'_2 \otimes \omega'_1 \omega'_2 f_1 \otimes f_1 \omega'_2 \otimes f_2 \\
& + \omega'_1{}^2 \omega'_2 f_1 \otimes f_1 \omega'_1 \omega'_2 \otimes f_1 \omega'_2 \otimes f_2 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_1 f_1 \omega'_2 \otimes f_1 \omega'_2 \otimes f_2 \\
& + f_1 \omega'_1{}^2 \omega'_2 \otimes f_1 \omega'_1 \omega'_2 \otimes f_2 \omega'_1 \otimes f_1 + \omega'_1 f_1 \omega'_1 \omega'_2 \otimes f_1 \omega'_1 \omega'_2 \otimes f_2 \omega'_1 \otimes f_1 \\
& + f_1 \omega'_1{}^2 \omega'_2 \otimes \omega'_1 \omega'_2 f_1 \otimes \omega'_1 f_2 \otimes f_1 + \omega'_1 f_1 \omega'_1 \omega'_2 \otimes \omega'_1 \omega'_2 f_1 \otimes \omega'_1 f_2 \otimes f_1 \\
& + \omega'_1{}^2 \omega'_2 f_1 \otimes f_1 \omega'_1 \omega'_2 \otimes \omega'_1 f_2 \otimes f_1 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_1 f_1 \omega'_2 \otimes \omega'_1 f_2 \otimes f_1 \\
& + f_1 \omega'_1{}^2 \omega'_2 \otimes \omega'_1 f_2 \omega'_1 \otimes f_1 \omega'_1 \otimes f_1 + f_1 \omega'_1{}^2 \omega'_2 \otimes \omega'_1 f_2 \omega'_1 \otimes \omega'_1 f_1 \otimes f_1 \\
& + \omega'_1 f_1 \omega'_1 \omega'_2 \otimes \omega'_1 f_2 \omega'_1 \otimes f_1 \omega'_1 \otimes f_1 + \omega'_1 f_1 \omega'_1 \omega'_2 \otimes \omega'_1 f_2 \omega'_1 \otimes \omega'_1 f_1 \otimes f_1 \\
& + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_1{}^2 f_2 \otimes f_1 \omega'_1 \otimes f_1 + \omega'_1{}^2 \omega'_2 f_1 \otimes \omega'_1{}^2 f_2 \otimes \omega'_1 f_1 \otimes f_1 \\
& + \omega'_1{}^2 f_2 \omega'_1 \otimes f_1 \omega'_1{}^2 \otimes f_1 \omega'_1 \otimes f_1 + \omega'_1{}^2 f_2 \omega'_1 \otimes \omega'_1 f_1 \omega'_1 \otimes f_1 \omega'_1 \otimes f_1 \\
& + \omega'_1{}^2 f_2 \omega'_1 \otimes f_1 \omega'_1{}^2 \otimes \omega'_1 f_1 \otimes f_1 + \omega'_1{}^2 f_2 \omega'_1 \otimes \omega'_1 f_1 \omega'_1 \otimes \omega'_1 f_1 \otimes f_1 \\
& + \omega'_1{}^2 f_2 \omega'_1 \otimes \omega'_1{}^2 f_1 \otimes f_1 \omega'_1 \otimes f_1 + \omega'_1{}^2 f_2 \omega'_1 \otimes \omega'_1{}^2 f_1 \otimes \omega'_1 f_1 \otimes f_1. \tag{4.3}
\end{aligned}$$

The relevant terms of  $\Delta(X)$  for  $X = f_1 f_2 f_1^2$  in (6C) are as follows:

$$\begin{aligned}
& f_1 \omega_1'^2 \omega_2' \otimes \omega_2' f_1 \omega_1' \otimes \omega_2' f_1 \otimes f_2 + f_1 \omega_1'^2 \omega_2' \otimes \omega_1' \omega_2' f_1 \otimes \omega_2' f_1 \otimes f_2 \\
& + \omega_1' \omega_2' f_1 \omega_1' \otimes f_1 \omega_1' \omega_2' \otimes \omega_2' f_1 \otimes f_2 + \omega_1'^2 \omega_2' f_1 \otimes f_1 \omega_1' \omega_2' \otimes \omega_2' f_1 \otimes f_2 \\
& + \omega_1' \omega_2' f_1 \omega_1' \otimes \omega_1' \omega_2' f_1 \otimes f_1 \omega_2' \otimes f_2 + \omega_1'^2 \omega_2' f_1 \otimes \omega_1' \omega_2' f_1 \otimes f_1 \omega_2' \otimes f_2 \\
& + f_1 \omega_1'^2 \omega_2' \otimes \omega_2' f_1 \omega_1' \otimes f_2 \omega_1' \otimes f_1 + f_1 \omega_1'^2 \omega_2' \otimes \omega_1' \omega_2' f_1 \otimes f_2 \omega_1' \otimes f_1 \\
& + \omega_1' \omega_2' f_1 \omega_1' \otimes f_1 \omega_1' \omega_2' \otimes f_2 \omega_1' \otimes f_1 + \omega_1'^2 \omega_2' f_1 \otimes f_1 \omega_1' \omega_2' \otimes f_2 \omega_1' \otimes f_1 \\
& + \omega_1' \omega_2' f_1 \omega_1' \otimes \omega_1' \omega_2' f_1 \otimes \omega_1' f_2 \otimes f_1 + \omega_1'^2 \omega_2' f_1 \otimes \omega_1' \omega_2' f_1 \otimes \omega_1' f_2 \otimes f_1 \\
& + f_1 \omega_1'^2 \omega_2' \otimes f_2 \omega_1'^2 \otimes f_1 \omega_1' \otimes f_1 + f_1 \omega_1'^2 \omega_2' \otimes f_2 \omega_1'^2 \otimes \omega_1' f_1 \otimes f_1 \\
& + \omega_1' \omega_2' f_1 \omega_1' \otimes \omega_1' f_2 \omega_1' \otimes f_1 \omega_1' \otimes f_1 + \omega_1'^2 \omega_2' f_1 \otimes \omega_1' f_2 \omega_1' \otimes f_1 \omega_1' \otimes f_1 \\
& + \omega_1' \omega_2' f_1 \omega_1' \otimes \omega_1' f_2 \omega_1' \otimes \omega_1' f_1 \otimes f_1 + \omega_1'^2 \omega_2' f_1 \otimes \omega_1' f_2 \omega_1' \otimes \omega_1' f_1 \otimes f_1 \\
& + \omega_1' f_2 \omega_1'^2 \otimes f_1 \omega_1'^2 \otimes f_1 \omega_1' \otimes f_1 + \omega_1' f_2 \omega_1'^2 \otimes f_1 \omega_1'^2 \otimes \omega_1' f_1 \otimes f_1 \\
& + \omega_1' f_2 \omega_1'^2 \otimes \omega_1' f_1 \omega_1' \otimes f_1 \omega_1' \otimes f_1 + \omega_1' f_2 \omega_1'^2 \otimes \omega_1'^2 f_1 \otimes f_1 \omega_1' \otimes f_1 \\
& + \omega_1' f_2 \omega_1'^2 \otimes \omega_1' f_1 \omega_1' \otimes \omega_1' f_1 \otimes f_1 + \omega_1' f_2 \omega_1'^2 \otimes \omega_1'^2 f_1 \otimes \omega_1' f_1 \otimes f_1.
\end{aligned} \tag{4.4}$$

The relevant terms of  $\Delta^{(2)}(f_2^2 f_1)$  in (7C) are as follows:

$$\begin{aligned}
& f_2 \omega_1' \omega_2' \otimes f_2 \omega_1' \otimes f_1 + \omega_2' f_2 \omega_1' \otimes f_2 \omega_1' \otimes f_1 \\
& + f_2 \omega_1' \omega_2' \otimes \omega_2' f_1 \otimes f_2 + \omega_2' f_2 \omega_1' \otimes \omega_2' f_1 \otimes f_2 \\
& + \omega_2'^2 f_1 \otimes f_2 \omega_2' \otimes f_2 + \omega_2'^2 f_1 \otimes \omega_2' f_2 \otimes f_2.
\end{aligned} \tag{4.5}$$

The relevant terms of  $\Delta^{(2)}(f_2 f_1 f_2)$  in (7C) are as follows:

$$\begin{aligned}
& f_2 \omega_1' \omega_2' \otimes \omega_1' f_2 \otimes f_1 + \omega_1' \omega_2' f_2 \otimes f_2 \omega_1' \otimes f_1 \\
& + f_2 \omega_1' \omega_2' \otimes f_1 \omega_2' \otimes f_2 + \omega_1' \omega_2' f_2 \otimes \omega_2' f_1 \otimes f_2 \\
& + \omega_2' f_1 \omega_2' \otimes f_2 \omega_2' \otimes f_2 + \omega_2' f_1 \omega_2' \otimes \omega_2' f_2 \otimes f_2.
\end{aligned} \tag{4.6}$$

### Acknowledgments

The study of the two-parameter quantum groups of types  $B$ ,  $C$ ,  $D$  was early initiated when NH visited l'Institut de Recherche Mathématique Avancée (IRMA – C.N.R.S), Strasbourg in the academic year 2000-2001, he would like to thank Prof. C. Kassel for his hospitality. YG would like to thank Professors M. Kashiwara and K. Saito for their extreme hospitality during his visit to RIMS of Kyoto University when part of this work was done. A special thanks go to Honglian Zhang for verifying Lemma 3.6.

## REFERENCES

1. M. Artin, W. Schelter, and J. Tate, *Quantum deformations of  $GL_n$* , Comm. Pure. Appl. Math **XLIV** (1991), 879–895.
2. G. Benkart, *Down-up algebras and Witten’s deformations of the universal enveloping algebra of  $\mathfrak{sl}_2$* ; *Recent Progress in Algebra*, Contemp. Math., vol 224, Amer. Math. Soc., 1998, pp.29–45.
3. G. Benkart and S. Witherspoon, *Two-parameter quantum groups and Drinfel’d doubles*, Algebr. Represent. Theory **7** (2004), 261–286.
4. ———, *Representations of two-parameter quantum groups and Schur-Weyl duality*, Hopf algebras, 65–92, Lecture Notes in Pure and Appl. Math., 237, Dekker, New York, 2004.
5. ———, *Restricted two-parameter quantum groups*, Fields Institute Communications, “Representations of Finite Dimensional Algebras and Related Topics in Lie Theory and Geometry”, Vol. 40, 2004, pp. 293–318.
6. W. Chin and I.M. Musson, *Multiparameter quantum enveloping algebras*, J. Pure & Appl. Algebra **107** (1996), 171–191.
7. V.K. Dobrev and P. Parashar, *Duality for multiparametric quantum  $GL(n)$* , J. Phys. A: Math. Gen. **26** (1993), 6991–7002.
8. V.G. Drinfeld, *Quantum groups*, in “Proceedings ICM”, Berkeley, Amer. Math. Soc. (1987), 798–820.
9. J. Du, B. Parshall and J.P. Wang, *Two-parameter quantum linear groups and the hyperbolic invariance of  $q$ -Schur algebras*, J. of London Math. Soc. **44** (1992), 420–436.
10. J.C. Jantzen, *Lectures on Quantum Groups*, vol. 6, Graduate Studies in Math. Amer. Math. Soc. Providence, 1996.
11. N.H. Jing, *Quantum groups with two parameters; Deformation Theory and Quantum Groups with Applications to Mathematical Physics (Amherst, MA, 1990)*, Contemp. Math., vol. 134, Amer. Math. Soc., 1992, pp. 129–138.
12. A. Joseph, *Quantum Groups and Their Primitive Ideals*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag, Berlin, 1995.
13. A. Klimyk and K. Schmüdgen, *Quantum Groups and Their Representations*, Springer-Verlag Berlin Heidelberg New York, 1997.
14. P.P. Kulish, *A two-parameter quantum group and gauge transformations (in Russian)*, Zap. Nauch. Semin. LOMI **180** (1990), 89–93.
15. G. Lusztig, *Introduction to Quantum Groups*, Birkhäuser Boston, 1993..
16. ———, *Quantum deformations of certain simple modules over enveloping algebras*, Adv. Math. **70** (1988), 237–249.
17. ———, *Quantum groups at roots of 1*, Geom. Dedicata **35** (1990), 89–114.
18. N. Reshetikhin, *Multiparameter quantum groups and twisted quasitriangular Hopf algebras*, Lett. Math. Phys. **20** (1990), 331–335.
19. N.Yu. Reshetikhin, L.A. Takhtajan and L.D. Faddeev, *Quantization of Lie groups and Lie algebras*, Algebra and Anal. **1** (1989), (Leningrad Math. J. 1 [Engl. transl. 193–225 (1990)]), 178–206.
20. M. Rosso, *Finite dimensional representations of the quantum analog of the enveloping algebra of a complex simple Lie algebra*, Comm. Math. Phys. **117** (1988), 581–593.
21. A. Sudbery, *Consistent multiparameter quantization of  $GL(n)$* , J. Phys. A: Math. Gen (1990), L697–L704.
22. M. Takeuchi, *A two-parameter quantization of  $GL(n)$* , Proc. Japan Acad. **66 Ser A** (1990), 112–114.